Noncommutative Geometry

Raimar Wulkenhaar

Mathematisches Institut der Westfälischen Wilhelms-Universität

Münster, Germany



Gelfand-Naimark Serre-Swan Spectral triples Operators HH Algebra Openness Tentative charts Reconstruction

Topological spaces

topological space X: set of points together with collection T (topology) of open subsets

- $Y \subset X$ closed if $X \setminus Y$ open
- sufficies to define convergence of sequences
- Suffices to define continuity: *φ* : *X* → Y continuous if for every open *Z* ⊂ Y the pre-image *φ*⁻¹(*Z*) is open in *X*
- $\phi: X \to Y$ homeomorphism if bijective and both ϕ, ϕ^{-1} continuous
- X is compact if any open cover has a finite subcover;
 X is sequentially compact if any sequence has a convergent subsequence (equivalent for metric spaces)

X may carry different topologies which leads to different notions of convergence and continuity!

Hausdorff spaces

Hausdorff space X: topological space in which distinct points are separated by open neighbourhoods

- Iimit of a convergent sequence is unique
- 2 compact subsets are closed
- all metric spaces are Hausdorff

Hausdorff space *X* is locally compact if every point has a compact neighbourhood

a locally compact Hausdorff space can be embedded in a compact Hausdorff space which has only one extra point at infinity

Gelfand-Naimark Serre-Swan Spectral triples Operators HH Algebra Openness Tentative charts Reconstruction 00 00000000 000 0000 000 0000</t

C*-algebras

- algebra A: vector space over K + ring + compatibility we assume 1 ∈ A and K = C
- normed algebra: || || : A → ℝ satisfying norm axioms of vector spaces, ||ab|| ≤ ||a|| ||b|| and ||1|| = 1
- Banach algebra: completeness, i.e. Cauchy sequences in *A* have a limit in *A*
- involution: $(a+\lambda b)^* = a^* + \overline{\lambda} b^*$, $(ab)^* = b^* a^*$, $(a^*)^* = a$
- C*-algebra: Banach *-algebra with $||a^*a|| = ||a||^2$

The C^* -property is very restrictive:

- If $\| \|$ unique: $\|a\|^2 = \sup\{|\lambda| : a^*a \lambda 1 \text{ not invertible in } A\}$
- **2** $\phi : \mathbf{A} \to \mathbf{B}$ isomorphism $\Rightarrow \|\phi(\mathbf{a})\| = \|\mathbf{a}\|$
- any C*-algebra is *-isomorphic to a norm-closed subalgebra of B(H) (bounded operators on Hilbert space)

Standard example: continuous functions

Operators

Spectral triples

A = C(X) continuous functions on compact Hausdorff space X with $||f|| := \sup_{x \in X} |f(x)|$, $(f^*)(x) := \overline{f(x)}$

0000000 000 00000

Algebra

Openness

Tentative charts

Reconstruction

Theorem. C(X) is a C^* -algebra

• norm-closed: Cauchy sequence $\{f_k\}$ in C(X) defines pointwise limit function f; it follows $||f - f_k|| \rightarrow 0$

 $\frac{\epsilon}{3}$ -trick proves continuity of f

2 C*-property:
$$\exists p \in X$$
 with $||f|| = \sup_{x \in X} |f(x)| = |f(p)|$

 $||f^*f|| = \sup_{x \in X} |f(x)|^2 = |f(p)|^2 = ||f||^2$

For X is locally compact: continuous functions vanishing at ∞ $C_0(X) = \{ f \in C(X) : \forall \epsilon > 0 \exists K \subset X \text{ compact with} |f(x)| < \epsilon \forall x \in X \setminus K \}$

Gelfand-Naimark

00000000000

Serre-Swan

Standard example: bounded operators

Spectral triples Operators

 \mathcal{H} complex Hilbert space, $\mathcal{B}(\mathcal{H})$ algebra of bounded linear operators on \mathcal{H} with $\|a\| := \sup_{x \in \mathbb{H}, \|x\|=1} \|ax\|$

Algebra

0000000 000 00000

Openness

Tentative charts

Reconstruction

adjoint operator $\langle a^*x, y \rangle = \langle x, ay \rangle$ from Riesz representation theorem, with $||a^*|| = ||a||$

Theorem. $\mathcal{B}(\mathcal{H})$ is a C^* -algebra

o norm-closed: same argument as for C(X)

 $0 \leq \|ax\|^2 = \langle ax, ax \rangle = \langle a^*ax, x \rangle \leq \|a^*ax\| \|x\| \leq \|a^*a\| \|x\|^2$

Example: *H* = L²(X, μ), then a C*-subalgebra of B(*H*) is L[∞](X, μ) with ||[f]|| = ess sup_μ{|f(x)| : x ∈ X} (in fact even a von Neumann algebra)

if
$$(X,\mu)$$
 a manifold: $\mathcal{A}=C^\infty(X)\subset C(X)\subset L^\infty(X,\mu)=\mathcal{A}''$

Gelfand-Naimark

00000000000

Serre-Swan

Gelfand-Naimark	Serre-Swan	Spectral triples	Operators	HH	Algebra	Openness	Tentative charts	Reconstruction
00000000000								

Spectrum

- A Banach algebra, $1 \in A$
 - spectrum of a ∈ A: sp(a) = {λ ∈ C : a − λ1 not invertible in A} resolvent set R(a) = C \ sp(a)
 - spectral radius $r(a) = \sup\{|\lambda| : \lambda \in sp(a)\}$

spectral radius formula: $r(a) = \lim_{n \to \infty} ||a^n||^{\frac{1}{n}} \le ||a||$ (equality for normal elements in *C**-algebra)

• R(a) open

resolvent function $z \mapsto (a - z1)^{-1}$ holomorphic on R(a)sp(a) nonempty and compact in \mathbb{C}

Characters

A - commutative C*-algebra

character: non-zero *-algebra homomorphism $\chi : \mathbf{A} \to \mathbb{C}$

 $\chi(1) = 1, ||\chi|| = 1$

$X = \operatorname{Spec}(A) = \{\chi : \chi \text{ character on } A\}$

- view X as subspace of unit ball A'₁ in dual space A'
- A' = {f : A → C linear} equipped with weak-*-topology open subsets generated by {â⁻¹(U) : U ⊂ C open, a ∈ A} with â(f) = f(a)
- weak-*-topology on A' separates points A'₁ ⊂ A' is compact (Banach-Alaoglu) X = Spec(A) ⊂ A'₁ closed

X with weak-*-topology is compact Hausdorff space

attention: A'_1 not compact in norm topology

The Gelfand-Naimark theorem

Gelfand transformation: $\rho : A \ni a \mapsto \hat{a} \in C(\text{Spec}(A))$, with $\hat{a}(\chi) := \chi(a)$, for commutative Banach algebra A

Theorem

Let *A* be a commutative *C*^{*}-algebra and *X* = Spec(*A*). Then $\rho : A \rightarrow C(X)$ is an isometric isomorphism.

We outline proof of $\|\hat{a}\| = \|a\|$ (injectivity). Surjectivity follows from Stone-Weierstraß.

•
$$|\hat{\boldsymbol{a}}(\chi)| = |\chi(\boldsymbol{a})| \le ||\chi|| ||\boldsymbol{a}|| = ||\boldsymbol{a}||$$

 $||\hat{\boldsymbol{a}}|| = \sup_{\chi \in X} |\hat{\boldsymbol{a}}(\chi)| \le ||\boldsymbol{a}||$

 If we can prove that for every λ ∈ sp(a) there is a character χ ∈ X with χ(a) = λ, then

$$\|a\|^2 = \|a^*a\| = r(a^*a) = \chi(a^*a) = \chi(a^*)\chi(a) = |\chi(a)|^2 = |\hat{a}(\chi)|^2$$

so $\|a\| \le \|\hat{a}\|$

I maximal if not contained in other ideal $\neq A$

• ker χ is ideal (clear) and maximal:

• $a \in I \setminus \ker \chi \Rightarrow \chi(a)$ invertible , $\exists b \in A: \chi(b)\chi(a) = 1$

• $ab \in I$ and $ab - 1 \in \ker \chi \subset I$, so $1 \in I$, contradiction

every maximal ideal *I* in commut. alg. A is of that type:

- A/I is a field; otherwise ∃0 ≠ b ∈ A/I not invertible.
 Then Ab ≇ 1 is an ideal, and I + Ab is a bigger ideal (b ∉ I)
- This field is C: Take 0 ≠ b ∈ A/I and λ ∈ sp(b), then b − λ1 not invertible, so b = λ1.
- Composition $A \to A/I \to \mathbb{C}$ defines character χ_I , with $I = \ker \chi_I$

Take $\lambda \in sp(a)$. Then $A(a - \lambda 1) \not\supseteq 1$ is ideal contained in maximal ideal ker χ .

$$1 \in A \Rightarrow \chi(a) = \lambda$$

The evaluation map

Given compact Hausdorff space X, then Y = Spec(C(X)) homeomorphic to X, because:

- evaluation map $\epsilon_x : C(X) \to \mathbb{C}, \quad \epsilon_x(f) = f(x)$ character
- **2** $\epsilon : X \to Y$ injective because C(X) separates points
- every character is of this type:
 - Take $\chi \in Spec(C(X))$ with $I = \ker \chi$ different from all $\ker \epsilon_x$.
 - For every $x \in X$ there is an $a_x \in I \subset C(X)$ with $a_x(x) \neq 0$.
 - a_x continuous $\Rightarrow a_x$ non-zero on neighbourhood of x
 - X compact $\Rightarrow \exists$ finitely many $x_1, \dots, x_n \in X$ with $a = \sum_{i=1}^n |a_{x_i}|^2 \in I$ non-zero on X
 - $\exists a^{-1} \in A$ with $1 = a^{-1}a \in I$, contradiction
- Sontinuity of ϵ, ϵ^{-1} by definition of weak-*-topology

Gelfand-Naimark	Serre-Swan	Spectral triples	Operators	ΗH	Algebra	Openness	Tentative charts	Reconstruction
0000000000								

A dictionary

compact Hausdorff space	commutative C*-algebra
measure space (X, μ)	commut. von Neumann algebra
group	commutative Hopf algebra
vector bundle over X	finitely generated projective mod-
	ule over $C(X)$
vector field	derivation
K-theory	K-theory
de Rham complex	Hochschild cohomology
de Rham cohomology	cyclic homology
differentiable manifold M	commutative spectral triple
diffeomorphism of M	automorphism of commutative
	spectral triple
(real/infinitesimal) variable	linear (selfadjoint/compact) oper-
	ator on Hilbert space
integral	trace

Vector bundles

Let *X* be a compact Hausdorff space. A locally-trivial vector bundle over *X* consists of a topological space *E* and a continuous surjection $p : E \to X$ such that

- For every $x \in X$, the fibre $E_X = p^{-1}(x) \subset E$ is a finite-dimensional complex vector space
- Solution For every *x* ∈ *X* there is a neighbourhood *U* and *m* ∈ N such that $p^{-1}(U)$ is homeomorphic to $\mathbb{C}^m \times U$.

A global section of $E \xrightarrow{p} X$ is a continuous mapping $s : X \to E$ with $p \circ s = id_X$.

We let $\Gamma(E, S)$ be the vector space of sections of $E \xrightarrow{p} X$. It becomes a module over C(X) by (sf)(x) := s(x)f(x) for $f \in C(X)$.

- A C(X)-module *E* is finitely generated if there exists a family η₁,..., η_k ∈ *E* such that every η ∈ *E* can be (not uniquely) represented as η = ∑_{i=1}^k η_if_i for f_i ∈ C(X).
- A C(X)-module *E* is free if it is homeomorphic to C^m ⊗ C(X) for some m ∈ N. A family of generators is e.g. (e_i ⊗ 1)_{i=1,...,m} for any basis (e_i) of C^m.
- The Whitney sum of vector bundles *E*₁, *E*₂ over *X* is the vector bundle

$$E_1 \oplus E_2 = \{(e_1, e_2) \in E_1 \times E_2 : p_1(e_1) = p_2(e_2)\}$$

- The corresponding direct sum $\mathcal{E}_1 \oplus \mathcal{E}_2$ of C(X)-modules $\mathcal{E}_1, \mathcal{E}_2$ becomes a C(X)-module by $(\eta_1, \eta_2)f := (\eta_1 f, \eta_2 f)$.
- A C(X)-Modul *E* is projective if there is another C(X)-module *E*' such that *E* ⊕ *E*' is a free module.

The Serre-Swan theorem

Theorem (Swan, 1962)

Let X be a compact Hausdorff space. A C(X)-module \mathcal{E} is isomorphic to a module $\Gamma(E, X)$ for a locally-trivial vector bundle $E \xrightarrow{p} X$ if and only iff \mathcal{E} is finitely generated and projective.

• The original article

R. G. Swan, "Vector Bundles and Projective Modules," Trans. Am. Math. Soc. **105** (1962) 264–277

is a good reference

 The theorem generalises to e.g. C[∞](X)-modules and for some measure space to L[∞]-modules

Let $E \xrightarrow{p} X$ be a locally-trivial vector bundle.

- For $x \in X$ there is a neighbourhood U such that $p^{-1}(U) \simeq \mathbb{C}^m \times U$.
- By surjectivity there exist $s_{x,1}, \ldots, s_{x,m} \in \Gamma(E, X)$ such that $\{pr_1(s_{x,1}(x)), \ldots, pr_1(s_{x,m}(x))\}$ form a basis of \mathbb{C}^m .
- By continuity there is a neighbourhood U_x ⊂ U of x such that {pr₁(s_{x,1}(y)),..., pr₁(s_{x,m}(y))} are also linearly independent for every y ∈ U_x.

This means that the family $\{s_{x,1}, \ldots, s_{x,m}\}$ generates $p^{-1}(U_x)$.

X is compact, i.e. covered by finitely many U_{x_1}, \ldots, U_{x_k} .

Thus finitely many $s_1(x), \ldots, s_n(x)$ generate $\Gamma(E, X)$

 Gelfand-Naimark
 Serre-Swan
 Spectral triples
 Operators
 HH
 Algebra
 Openness
 Tentative charts
 Reconstruction

 000000000
 000
 000
 000
 000
 000
 000
 000
 000
 0000
 000
 00000000
 000
 0000
 000
 00000000
 000
 0000
 0000
 0000
 0000
 0000
 0000
 0000
 0000
 0000
 0000
 0000
 0000
 0000
 0000
 0000
 0000
 0000
 0000
 0000
 0000
 0000
 0000
 0000
 0000
 0000
 0000
 0000
 0000
 0000
 0000
 0000
 0000
 0000
 0000
 0000
 0000
 0000
 0000
 0000
 0000
 0000
 0000
 0000
 0000
 0000
 0000
 0000
 0000
 0000
 0000
 0000
 0000
 0000
 0000
 0000
 0000
 0000
 0000
 0000
 0000
 0000
 0000
 0000</

previous page: finitely many $s_1(x), \ldots, s_n(x)$ generate $\Gamma(E, X)$

- Consider the free C(X)-module $\mathcal{F} = \mathbb{C}^n \otimes C(X) = \Gamma(\mathbb{C}^n \times X, X)$ with *n* generators b_1, \ldots, b_n .
- Define a mapping e : F → E by e(b_i) := s_i and extension by the module structure. This mapping is surjective and continuous.
- The dimension of the image of *e* is locally constant. Thus, the dimension of the kernel of *e* is locally constant, too.
- Locally constant dimension is sufficient to reconstruct a vector bundle (Proposition 1 in Swan's paper).
- Both $im(e) = \mathcal{E}$ and $ker(e) =: \mathcal{E}'$ are C(X)-modules, and $\mathcal{F} = \mathcal{E} \oplus \mathcal{E}'$.

Hence, $\ensuremath{\mathcal{E}}$ is projective.

Proof (⇐)

Let \mathcal{E} be a finitely generated projective C(X)-module, $\mathcal{E} \oplus \mathcal{E}' = \mathcal{F}$ and $\xi = (\eta, \eta') \in \mathcal{F} = \mathbb{C}^n \otimes C(X)$.

• Define $\mathbf{e}: \mathcal{F} \to \mathcal{F}$ by $\mathbf{e}\xi = (\eta, \mathbf{0})$.

Then $e^2 = e$, $(e\xi)f = e(\xi f)$ and $\mathcal{E} \simeq im(e)$, $\mathcal{E}' \simeq ker(e)$.

- For $x \in X$ consider maximal ideal $I_x = \{f \in C(X) : f(x) = 0\}$ in C(X).
- \$\mathcal{F}I_x\$ consists of elements of \$\mathcal{F}\$ vanishing in \$x\$
 \$\mathcal{F}/\mathcal{F}I_x\$ equivalence classes with same value in \$x\$
- Evaluation $\xi \mapsto \xi(x)$ defines isomorphism $\mathcal{F}/\mathcal{F}I_x \simeq F_x \simeq \mathbb{C}^n$.
- $(e\xi)f = e(\xi f)$

The evaluation $e\xi \mapsto (e\xi)(x)$ gives an isomorphism $e\mathcal{F}/e\mathcal{F}I_x \simeq E_x$ with a subspace $E_x \subset \mathbb{C}^m$.

If we can show that the dimension of E_x is locally constant, then the fibres E_x define a locally-trivial vector bundle.

- Let m := dim(E_x). There exist m linearly independent continuous sections s₁,..., s_m of Cⁿ × X, i.e. s_i ∈ F, with (es_i)(x) = s_i(x).
- By continuity, the (es_i) are linearly independent in a neighbourhood U_x of x, i.e. dim(E_y) ≥ m for all y ∈ U_x.
- Same argument for 1 e gives complementary fibre space E'_x of dimension n m. Consequently, $\dim(E'_y) \ge n m$.
- Total dimension n is constant, thus dim(E_x) = m locally constant.

Gelfand-Naimark	Serre-Swan	Spectral triples	Operators	ΗН	Algebra	Openness	Tentative charts	Reconstruction
		•00						

Spectral triples

Definition (A. Connes, 1996)

 $(\mathcal{A}, \mathcal{H}, D)$ – commutative spectral triple, i.e. \mathcal{H} a Hilbert space, \mathcal{A} a commutative involutive unital algebra represented in \mathcal{H} , D a selfadjoint operator in \mathcal{H} with compact resolvent, p an integer.

O Dimension: k^{th} characteristic value of resolvent of *D* is $\mathcal{O}(k^{-\frac{1}{p}})$

2 Order one:
$$[[D, f], g] = 0 \quad \forall f, g \in A$$

Segularity: *f* and [*D*, *f*] belong to the domain of δ^m , for any $f \in \mathcal{A}$ and $m \in \mathbb{Z}$, where $\delta T := [|D|, T]$

Orientability: ∃ Hochschild *p*-cycle *c* ∈ Z_p(A, A) s.t. π_D(*c*) = 1 for *p* odd, π_D(*c*) = γ for *p* even with γ = γ*, γ² = 1, γD = −Dγ

Siniteness and absolute continuity: $\mathcal{H}_{\infty} := \bigcap_{m} \operatorname{dom}(D^{m}) \subset \mathcal{H}$ is finitely generated projective \mathcal{A} -module, $\mathcal{H}_{\infty} = e\mathcal{A}^{n}$, with $e = e^{*} = e^{2} \in M_{n}(\mathcal{A})$. Hermitian structure $(\xi | a\eta) = \sum_{i=1}^{n} a\xi_{i}^{*}\eta_{i} \in \mathcal{A}$ satisfies $\langle \xi, \eta \rangle = f(\xi | \eta) |D|^{-p}$

Reconstruction theorem

Theorem (A. Connes, 2008)

Let $(\mathcal{A}, \mathcal{D}, \mathcal{H})$ be a spectral triple, \mathcal{A} commutative and unital. Let the conditions (1)–(5) be realised in stonger form:

 $\textcircled{ \textbf{ 0 } All } T \in \operatorname{End}_{\mathcal{A}}(\mathcal{H}_{\infty}) \text{ are regular }$

 $\begin{array}{l} \hline \textbf{O} \quad \textit{The Hochschild cycle is antisymmetric,} \\ \textbf{c} = \sum_{\alpha} \textbf{a}_{\alpha}^{0} \otimes \sum_{\beta \in \textbf{S}_{p}} \epsilon(\beta) \textbf{a}_{\alpha}^{\beta(1)} \otimes \cdots \otimes \textbf{a}_{\alpha}^{\beta(p)} \end{array}$

Then there exists a compact oriented differentialble manifold X with $\mathcal{A} = \mathbf{C}^{\infty}(\mathbf{X})$.

Conversely, every compact oriented differentiable manifold arises in this way.

Second theorem: If in addition the multiplicity of \mathcal{A}'' in \mathcal{H} is $2^{p/2}$, then $\mathcal{A} = C^{\infty}(X)$ for a smooth oriented compact spin^{*c*}-manifold *X*. The stronger condition (1) is automatic.

Definition (smooth compact *p*-dimensional manifold)

... is a compact Hausdorff space X together with a system of local charts (U_{α}, s_{α}) such that

- the U_{α} are open in X and $X = \bigcup_{\alpha} U_{\alpha}$
- $s_{\alpha}: U_{\alpha} \to s_{\alpha}(U_{\alpha}) \subset \mathbb{R}^{p}$ is a homeomorphism. In particular, $s_{\alpha}(U_{\alpha})$ is open in \mathbb{R}^{p} and s_{α} is injective

•
$$\mathbf{s}_{\alpha} \circ \mathbf{s}_{\beta}^{-1} : \mathbf{s}_{\beta}(U_{\alpha} \cap U_{\beta}) \to \mathbf{s}_{\alpha}(U_{\alpha} \cap U_{\beta})$$
 is smooth

Strategy:

- norm-completion of A is unital commutative C*-algebra
 A = C(X) for compact Hausdorff space X = Spec(A)
- build tentative charts (up to injectivity) from c
- prove that there exists restriction to subsets where s_α is injective (very complicated)



Unbounded operators

From (1), *D* is unbounded linear operator on \mathcal{H} for p > 0.

- $\operatorname{dom}(T) \subset \mathcal{H} \operatorname{domain}$
- $\Gamma(T) = \{(\phi, T\phi) \in \mathcal{H} \times \mathcal{H} : \phi \in \operatorname{dom}(T)\} \operatorname{graph}$

• T closed if $\Gamma(T)$ closed in $\mathcal{H} \times \mathcal{H}$

An extension of *T* is an operator T_1 with dom $(T_1) \supset$ dom(T) and $T_1\phi = T\phi$ for $\phi \in$ dom(T).

- T is closable if a closed extension exists.
- The smallest closed extension is the closure \overline{T} .

 $T : \operatorname{dom}(T) \to \mathcal{H} -$ linear densely defined operator

• dom(T^*) = { $\phi \in \mathcal{H}$: $\exists \eta \in \mathcal{H} \text{ with } \langle T\psi, \phi \rangle = \langle \psi, \eta \rangle$ for all $\psi \in \text{dom}(T)$ }

Then $T^*\phi := \eta$

• T^* always closed, T closable if dom $(T^*) \subset \mathcal{H}$ dense, with $\overline{T} = T^{**}$

Definition

A linear densely defined operator T is

- symmetric if dom(T) ⊂ dom(T*) and Tφ = T*φ for all φ ∈ dom(T)
- selfadjoint if T* = T, i.e. dom(T) = dom(T*) and T symmetric
- essentially selfadjoint if T is symmetric and \overline{T} selfadjoint

For closed operators one can define a spectral theory.

• resolvent set

$$R(T) = \{\lambda \in \mathbb{C} : \lambda 1 - T : \operatorname{dom}(T) \to \mathcal{H} \text{ bijective} \\ \text{and } (\lambda 1 - T)^{-1} \text{ bounded} \}$$

spectrum sp(T) = C \ R(T)
 sp(T) is closed, in general non-compact, possibly empty

If T is selfadjoint, then $sp(T) \subset \mathbb{R}$. Further:

- Cayley transformation $T \mapsto U_T := (i1 T)(-i1 T)^{-1}$ to unitary U_T with $1 U_T$ injective
- spectral theorem $T = \int_{\mathbb{R}} \lambda \ dE_{\lambda}$ for some spectral measure dE_{λ}

Polar decomposition T = F|T| for closed densely defined *T*:

- |T| positive and selfadjoint, dom(|T|) = dom(T), ker |T| = ker T
- $F: (\ker T)^{\perp} \to \overline{im(T)}$ partial isometry

Compact operators

Let $\mathcal{K} \subset \mathcal{B}(\mathcal{H})$ be the ideal of compact operators on \mathcal{H} and $\mathcal{T} \in \mathcal{K}$.

- There is a null sequence of eigenvalues $\mu_i > 0$ of |T|
- The eigenspace $E_i = \text{ker}(\mu_i 1 |T|)$ is finite-dimensional
- $sp(|T|) = {\mu_i}_{i \in \mathbb{N}} \cup {0}$

 $s_k(T) = \inf\{\|T|_{E^{\perp}}\| : \dim(E) = k\}$ – characteristic value

 $s_k(T) - k^{\text{th}}$ eigenvalue $\mu_k(|T|)$ if arrangend decreasingly and with multiplicity

Schatten ideal \mathcal{L}^p

$$\mathcal{L}^{p} := \left\{ T \in \mathcal{K} : \|T\|_{p} := \left(\sum_{k=0}^{\infty} (s_{k}(T))^{p} \right)^{\frac{1}{p}} < \infty \right\}$$

All $(\mathcal{L}^{p}, || ||_{p})$ are Banach spaces.

Gelfand-Naimark	Serre-Swan	Spectral triples	Operators	ΗH	Algebra	Openness	Tentative charts	Reconstruction
			00000000					

- \mathcal{L}^1 trace class $\operatorname{Tr}(T) = \sum_n \langle \psi_n, T\psi_n \rangle$ for $T \in \mathcal{L}^1$ and $\{\psi_n\}$ ONB
- \mathcal{L}^2 Hilbert-Schmidt class

•
$$\mathcal{L}^{\infty} = \mathcal{K}$$
 with $\|T\|_{\infty} = \|T\|$

Inequalities

- $\operatorname{Tr}(T) \leq \operatorname{Tr}(|T|) = ||T||_1$ for $T \in \mathcal{L}^1$
- Tr(TS) = Tr(ST) if $TS, ST \in \mathcal{L}^1$
- Hölder: $||TS||_1 \le ||T||_p ||S||_q$ for $T \in \mathcal{L}^p$, $S \in \mathcal{L}^q$ with $\frac{1}{p} + \frac{1}{q} = 1$
- $\mathcal{L}^{p} \subset \mathcal{L}^{r}$ if $p \leq r$

It turns out that Tr on \mathcal{L}^1 is not the right generalisation of the integral.

Dixmier ideal \mathcal{L}^{1+}

• partial sums $\sigma_n(T) = \sum_{k=0}^{n-1} s_k(T)$

Lemma

$$\sigma_n(T + S) \le \sigma_n(T) + \sigma_n(S) \le \sigma_{2n}(T + S)$$
 for $0 \le T, S \in \mathcal{K}$

Proof: first \leq from norm, second from inequalities

$$\begin{split} \|TP_E\|_1 + \|SP_F\|_1 &= \operatorname{Tr}(P_E TP_E) + \operatorname{Tr}(P_F SP_F) \\ &\leq \operatorname{Tr}(P_{E+F} TP_{E+F}) + \operatorname{Tr}(P_{E+F} SP_{E+F}) \\ &= \operatorname{Tr}(P_{E+F} (T+S)P_{E+F}) \end{split}$$

Definition (Dixmier-Ideal $\mathcal{L}^{1+} \subset \mathcal{K}$)

$$\mathcal{L}^{1+} := \left\{ T \in \mathcal{K} : \|T\|_{1+} := \sup_{n \ge 2} \frac{\sigma_n(T)}{\log n} < \infty \right\}$$

•
$$\mathcal{L}^1 \subset \mathcal{L}^{1+} \subset \mathcal{L}^p$$
 for any $p > 1$

If $\left(\frac{\sigma_n(T)}{\log n}\right)_{n\geq 2}$ convergent, then $T\in \mathcal{L}^{1+}$ is called measurable

Definition (noncommutative integral)

$$\int T := \lim_{n \to \infty} \frac{\sigma_n(T)}{\log n}$$

- \oint additive on positive elements, linear by extension
- $\sigma_n(UTU^*) = \sigma_n(T)$:

 \oint is trace on subspace of measurable elements of \mathcal{L}^{1+}

- \oint vanishes on \mathcal{L}^1
- $\bullet\,$ can be (not uniquely) generalised to Dixmier trace on \mathcal{L}^{1+}

Example

$$\Delta = -\sum_{\mu=1}^{p} rac{\partial^2}{\partial x_{\mu}^2}$$
 Laplace operator on \mathbb{T}^{p}

• $sp(\Delta) = \{ \|k\|^2 : k \in \mathbb{Z}^p \}$

• Replace Δ by 1 on ker $\Delta = \mathbb{C} \Rightarrow \tilde{\Delta}^{-\frac{q}{2}}$ compact for q > 0

• Determine $s_{n(||k||)}(\tilde{\Delta}^{-\frac{q}{2}}) = ||k||^{-q}$ asymptotically: $n(||k||) = #(\text{lattice points in ball of radius } ||k|| \text{ in } \mathbb{R}^p)$

$$= V_{\rho} \|k\|^{\rho}$$
 with $V_{\rho} = \frac{\pi^{\frac{1}{2}}}{\Gamma(\frac{\rho+2}{2})}$

•
$$s_n(\tilde{\Delta}^{-\frac{q}{2}}) = \left(\frac{n}{V_p}\right)^{-\frac{q}{p}}$$

•
$$\sigma_n(\tilde{\Delta}^{-\frac{q}{2}}) = \int_1^n du \left(\frac{V_p}{u}\right)^{\frac{q}{p}}, \quad \|\tilde{\Delta}^{-\frac{q}{2}}|_{1+} = \begin{cases} \infty & \text{for } q p \end{cases}$$
$$\int \tilde{\Delta}^{-\frac{p}{2}} = \lim_{n \to \infty} \frac{1}{\log n} \int_1^n du \, \frac{V_p}{u} = V_p$$

Hochschild homology

 \mathcal{A} unital algebra, \mathcal{M} bimodule over \mathcal{A} and $C_n(\mathcal{A}, \mathcal{M}) = \mathcal{M} \otimes \mathcal{A}^{\otimes n}$

Definition (Hochschild boundary operator)

$$b: C_n(\mathcal{A}, \mathcal{M}) \rightarrow C_{n-1}(\mathcal{A}, \mathcal{M}),$$

$$b(m \otimes a_1 \otimes \ldots \otimes a_n)$$

:= $ma_1 \otimes a_2 \otimes \cdots \otimes a_n$
+ $\sum_{i=1}^{n-1} (-1)^i m \otimes a_1 \otimes \cdots \otimes a_{i-1} \otimes a_i a_{i+1} \otimes a_{i+2} \otimes \cdots \otimes a_n$
+ $(-1)^n a_n m \otimes a_1 \otimes \cdots \otimes a_{n-1}$.

and b = 0 for n = 0

Gelfand-Naimark	Serre-Swan	Spectral triples	Operators	HH	Algebra	Openness	Tentative charts	Reconstruction
				000				

Definition

The homology of the chain complex $(C_*(\mathcal{A}, \mathcal{M}), b)$ is called Hochschild homology and denoted $HH_*(\mathcal{A}, \mathcal{M})$

Means: If $Z_n(\mathcal{A}, \mathcal{M}) \ni c_n$ is the subspace of Hochschild *n*-cycles $bc_n = 0$, then $HH_n(\mathcal{A}, \mathcal{M}) = Z_n(\mathcal{A}, \mathcal{M})/bC_{n+1}(\mathcal{A}, \mathcal{M})$

Theorem (Hochschild-Kostant-Rosenberg-Connes)

Let $A = C^{\infty}(X)$. Then $HH_*(C^{\infty}(X), C^{\infty}(X)) \simeq \Omega^*(C^{\infty}(X))$

remarkable fact: $HH_*(C^{\infty}(X), C^{\infty}(X))$ is local, only the diagonal in $(C^{\infty}(X))^{\otimes (n+1)} \simeq C^{\infty}(X \times \cdots \times X)$ contributes

- In spectral triple definition, the class [c] ∈ HH_p(A, A) of c ∈ Z_p(A, A) is the volume form (orientation class). It is local and nowhere vanishing because of γ² = 1, γ = π_D(c)
- c ∈ Z_p(A, A) and γ = 1 for p odd and γD = −Dγ for p even imply that |D|^{-p} ∈ L¹⁺ is measurable, i.e. f unambiguously defined

Gelfand-Naimark

Serre-Swan Spectral triples

Operators HH

I Algebra Open

Openness Tentative charts

arts Reconstruction

Characterisation of the algebra

Main Lemma

$$\mathcal{A} = \{ T \in \mathcal{A}'' : T \in \bigcap_{m > 0} \operatorname{dom}(\delta^m) \}$$

- *T* ∈ *A*" ⊃ *A* (both commutative) and *T* : *H*_∞ → *H*_∞ conclusion: *T* ∈ End_{*A*}(*H*_∞)
- \mathcal{A} is unital $\Rightarrow \operatorname{End}_{\mathcal{A}}(\mathcal{H}_{\infty}) = eM_n(\mathcal{A})e$ $T = eTe = (\alpha_{kl})$ with $\alpha_{kl} = (e\xi_k | Te\xi_l) \in \mathcal{A}$
- norm-completion of A is unital commutative C^* -algebra A = C(X) for X = Spec(A) a compact Hausdorff space
- define positive measure μ on X by

$$\mu(f) := \int f |D|^{-p} \qquad orall f \in \mathcal{A} = \mathcal{C}(X)$$

 $\mathcal{E} = eA^n = \mathcal{H}_{\infty} \otimes_{\mathcal{A}} A$ is finitely generated projective module

- Serre-Swan: ∃ a complex locally trivial vector bundle S → X such that E = Γ(X, S)
- Finiteness axiom $\Rightarrow \langle \xi, \eta \rangle = \mu((\xi|\eta))$ for $\xi, \eta \in e\mathcal{A}^n$
- $\mathcal{H}_{\infty} \subset \mathcal{H}$ dense from selfadjointness of D

completion of \mathcal{H}_∞ to \mathcal{H} is the same as completion with respect to μ

 $\Rightarrow \mathcal{H} = e(L^2(X,\mu))^n = L^2(X, \mathcal{S},\mu)$

action of f ∈ A" on H = L²(X, S, μ) is diagonal multiplication with f ∈ L[∞](X, μ)

$$T = e \operatorname{diag}(f, \ldots, f) \in eM_n(\mathcal{A}) \quad \Rightarrow \quad f \in \mathcal{A}$$

Gelfand-Naimark Serre-Swan Spectral triples

Operators 0000000 000 00000

Algebra Openness

Tentative charts

Reconstruction

• tr(e) = $\sum_{i=1}^{n} e_{ii} \in A \subset A$ is the continuous function which assigns to $\chi \in X$ the dimension of the fibre S_{χ} of S over χ . Thus, $\chi(tr(e)) \in \{1, \dots, n\}$ (0 can be excluded)

• Let $p_i \in A$ be the projection to the set of connected components X_i of X for which dim $(S_{\chi}) = j$. Then, $\sum_{i=1}^{n} p_i = 1$ and $\operatorname{tr}(e) = \sum_{i=1}^{n} j p_i \in \mathcal{A}$.

Reconstruction
$$p_k = \prod_{j \in \{1,...,n\} \setminus \{k\}} \frac{\operatorname{Tr}(e) - j\mathbf{1}}{k - j} \in \mathcal{A}$$

Definition (conditional expectation values)

Let $T = (T_{kl}) \in \operatorname{End}_{\mathcal{A}}(\mathcal{H}_{\infty}) = eM_{n}(\mathcal{A})e$.

$$E_{\mathcal{A}}(T) := \sum_{k=1}^{n} \frac{p_{k}}{k} \sum_{j=1}^{n} T_{jj} \in \mathcal{A}$$

examples:
$$E_{\mathcal{A}}(e) = 1$$
, $E_{\mathcal{A}}(e \operatorname{diag}(f, \dots, f)) = f$

Raimar Wulkenhaar (Münster) Noncommutative Geometry

Proposition

 \mathcal{A} is a Fréchet algebra, i.e. a complete locally convex algebra whose toplogy is defined by the submultiplicative norms $q_k(ab) \leq q_k(a)q_k(b)$,

$$q_k(a) = \|
ho_k(a)\|$$
, $ho_k(a) = egin{pmatrix} a & \delta(a) & \dots & \delta^k(a)/k! \ 0 & a & \dots & \dots \ \dots & \dots & a & \delta(a) \ 0 & \dots & 0 & a \end{pmatrix}$

Proof:

- If (a_n) Cauchy sequence in A w.r.t. any q_k, then (k = 0)
 a_n → T ∈ A ⊂ A"
- δ is closed by selfadjointness of *D*, thus $T \in \text{dom}(\delta)$.
- Inductively $T \in \text{dom}(\delta^m)$, thus $T \in \mathcal{A}$ from Main Lemma.

Proposition

 \mathcal{A} is a pre-*C**-algebra, i.e. closed under holomorphic functional calculus. This means: If $a \in \mathcal{A} \subset A$ and f holomorphic in neighbourhood of sp(a) (viewed from A), then

$$f(a) := rac{1}{2\pi \mathrm{i}} \oint_{c} dz rac{f(z)}{z 1 - a} \in \mathcal{A}$$

Proof:

- $z \notin sp(a) \Rightarrow (z1-a)^{-1} \in A \subset \mathcal{A}''$
- $(z1 a)^{-1}$ locally expanded in power series in $\zeta 1 a \in A$
- δ closed \Rightarrow $(z1 a)^{-1} \in \operatorname{dom}(\delta)$
- Inductively $(z1 a)^{-1} \in \text{dom}(\delta^m)$, thus $(z1 a)^{-1} \in \mathcal{A}$ from Main Lemma.
- convergence of Riemann integral in A by Fréchet

Consequence: $a \in A$ has the same spectrum in A and A $\Rightarrow X = Spec(A) = Spec(A)$

Gelfand-Naimark	Serre-Swan	Spectral triples	Operators	ΗH	Algebra	Openness	Tentative charts	Reconstruction
00000000000	0000000		00000000		00000	000	000000000	0000000000

Flows

Main Assumption

Let δ_0 be a continuous \star -derivation on the Fréchet pre- C^* -algebra \mathcal{A} . Then there exists a unique solution, depending continuously on $(t, a) \in \mathbb{R} \times \mathcal{A}$, of the differential equation

 $\partial_t y(t,a) = \delta_0(y(t,a)) , \quad y(0,a) = a$

Proposition

Let δ_0 be a continuous \star -derivation satisfying the Main Assumtion. Let $F_t(a) := y(t, a)$. Then $F_t \in Aut(\mathcal{A})$ is a one-parameter family of automorphisms, with smooth dependence on *t*.

Corollary

Let $\chi \in X = \text{Spec}(\mathcal{A})$ be a character, $F_t \in \text{Aut}(\mathcal{A})$ as before. Then $F_t^* \chi \in X$, where $F_t^* \chi(a) := \chi(F_t(a))$.

Main Data

- *p* derivations δ₁,..., δ_p satisfying the Main Assumption, with corresponding flows F¹_{t1},..., F^p_{tp} ∈ Aut(A),
- 2 *p* selfadjoint elements $a^1, \ldots, a^p \in A$,
- **③** a character $\chi \in X$.

Consider the map $\phi_{\chi} : \mathbb{R}^{p} \to \mathbb{R}^{p}$ with

$$\phi_{\chi}^{k}(t_{1},\ldots,t_{p})=\chi\left(F_{t_{1}}^{1}\circ\cdots\circ F_{t_{p}}^{p}(a^{k})\right)$$

It satisfies $(\partial_j \phi_{\chi}^k)(\mathbf{0}) = \chi(\delta_j \mathbf{a}^k)$.

Proposition

Given the Main Data (δ_i, a^k, χ) as before, with det $\chi(\delta_i a^k) \neq 0$.

There exists a neighbourhood Z ⊂ X = Spec(A) of χ and a neighbourhood W ⊂ ℝ^p of 0 such that, for any κ ∈ Z, the map

$$W
i t \mapsto \phi_{\kappa}(t) \in \mathbb{R}^{p}$$

is a diffeomorphism, depending continuously on κ , of W with a neighbourhood $Y_{\kappa} = \phi_{\kappa}(W)$ of $(\kappa(a^1), \ldots, \kappa(a^p)) \in \mathbb{R}^p$.

② The image of any open neighbourhood $U \subset X$ of χ under

$$U
i \kappa \mapsto (\kappa(a^1), \ldots, \kappa(a^p)) \in \mathbb{R}^p$$

contains an open neighbourhood of $(\chi(a^1), \ldots, \chi(a^p))$.

 Gelfand-Naimark
 Serre-Swan
 Spectral triples
 Operators
 HH
 Algebra
 Openness
 Tentative charts
 Reconstruction

 0000000000
 0000000
 000
 00000000
 000
 000
 00000000
 0000
 00000000
 000000000
 000000000
 000000000
 0000000000
 0000000000
 000000000
 000000000
 0000000000
 0000000000
 0000000000
 0000000000
 0000000000
 0000000000
 0000000000
 0000000000
 0000000000
 0000000000
 0000000000
 0000000000
 0000000000
 0000000000
 0000000000
 0000000000
 0000000000
 0000000000
 0000000000
 0000000000
 0000000000
 0000000000
 0000000000
 0000000000
 0000000000
 0000000000
 0000000000
 0000000000
 0000000000
 0000000000
 0000000000
 000000000
 0000000000
 0000000000
 0000000000
 0000000000
 0000000000
 00000000000
 0000000000
 0000000000
 00000000000
 0000000000
 0000000000
 00000000000
 000000000000
 000000000000000
 000000000000000000000000000000

Identifying the derivations

Lemma

One has a finite decomposition $[D, a] = \sum_{j=1}^{m} \delta_j(a)\gamma_j$ for all $a \in A$, where $\gamma_j \in \operatorname{End}_{\mathcal{A}}(\mathcal{H}_{\infty})$, and δ_j are *-derivations of \mathcal{A} of the form $\delta_j(a) = i(\eta_j, [D, a]\eta_j)$ for some $\eta_j \in \mathcal{H}_{\infty}$.

Proof

- $[D, a]\mathcal{H}_{\infty} \subset \mathcal{H}_{\infty}$ from $D[D, a]\xi = F\delta([D, a])\xi + F[D, a]|D|\xi$ etc. and regularity condition, $F = D|D|^{-1}$
- **order-one condition:** $b[D, a]\xi = [D, a]b\xi$ consequence: $[D, a] \in \text{End}_{\mathcal{A}}(\mathcal{H}_{\infty}) = eM_n(\mathcal{A})e$

$$[D, a] = \sum_{k,l} \alpha_{kl} \varepsilon_{kl}$$
 with $\alpha_{kl} = (e\xi_k, [D, a]e\xi_l) \in A$
with ξ_k units in A^n and ε_{kl} matrix units in $M_n(A)$

Sonsider A ∋ a → L_{kl}(a) := (eξ_k, [D, a]eξ_l) ∈ A. $L_{kl}(ab) = (eξ_k, [D, ab]eξ_l) = (eξ_k, [D, a]beξ_l) + (eξ_k, a[D, b]eξ_l) = (eξ_k, b[D, a]eξ_l) + (eξ_k, a[D, b]eξ_l) = (eξ_k, b[D, a]eξ_l) + (eξ_k, a[D, b]eξ_l) = b(eξ_k, [D, a]eξ_l) + a(eξ_k, [D, b]eξ_l) = L_{kl}(a)b + aL_{kl}(b)$

using order-one, $\mathcal A\text{-linearity}$ of Hermitian structure and commutativity of $\mathcal A$

Consequence: $[D, a] = \sum_{k,l} L_{kl}(a) \varepsilon_{kl}$ with $L_{kl} \in \text{Der}(\mathcal{A})$

achieve *-derivations by polarisation identity

$$2(\xi, T\eta) = (\xi + \eta, T(\xi + \eta)) - (\xi, T\xi) - (\eta, T\eta)$$
$$- i\{(\xi + i\eta, T(\xi + i\eta)) + i(\xi, T\xi) + i(i\eta, i\eta)\}$$
together with

 $(\xi, [D, a^*]\xi) = -(\xi, [D, a]^*\xi) = -([D, a]\xi, \xi) = -(\xi, [D, a]\xi)^*$

Gelfand-Naimark Serre-Swan Spectral triples Operators HH Algebra Openness Tentative charts coordooooo coord coord

Evaluation of γ

• Hochschild cycle

$$c = \sum_{\alpha} a_{\alpha}^{0} \otimes \sum_{\beta \in S_{p}} \epsilon(\beta) a_{\alpha}^{\beta(1)} \otimes \cdots \otimes a_{\alpha}^{\beta(p)} \text{ with } a_{\alpha}^{j} = (a_{\alpha}^{j})^{*}$$
• $\gamma = \pi_{D}(c) = \sum_{\alpha} a_{\alpha}^{0} \sum_{\beta \in S_{p}} \epsilon(\beta) [D, a_{\alpha}^{\beta(1)}] \cdots [D, a_{\alpha}^{\beta(p)}]$

$$= \sum_{\alpha} a_{\alpha}^{0} T_{\alpha}$$
• $T_{\alpha} = \sum_{j_{1}, \dots, j_{p}=1}^{m} \sum_{\beta \in S_{p}} \epsilon(\beta) \delta_{j_{1}}(a_{\alpha}^{\beta(1)}) \cdots \delta_{j_{p}}(a_{\alpha}^{\beta(p)}) \gamma_{j_{1}} \cdots \gamma_{j_{p}}$

$$T_{\alpha} = \sum_{\beta \in S_{p}} \epsilon(\beta) \sum_{j_{1}, \dots, j_{p}=1}^{m} \delta_{j_{1}}(\boldsymbol{a}_{\alpha}^{\beta(1)}) \cdots \delta_{j_{p}}(\boldsymbol{a}_{\alpha}^{\beta(p)}) \gamma_{j_{1}} \cdots \gamma_{j_{p}}$$

m

- if $j_k = j_l$ for $k \neq l$, then no contribution to the sum
- the remaining *j*-sum is split into the sum over the subsets
 F ⊂ {1,..., *m*} with |*F*| = *p* and the sum over the permutations *σ* of *F*:

$$T_{\alpha} = \sum_{\beta \in S_{p}} \epsilon(\beta) \sum_{1 \le j_{1} < \dots < j_{p} \le m} \sum_{\sigma \in S_{p}} \delta_{\sigma(j_{1})}(\boldsymbol{a}_{\alpha}^{\beta(1)}) \cdots \delta_{\sigma(j_{p})}(\boldsymbol{a}_{\alpha}^{\beta(p)}) \gamma_{\sigma(j_{1})} \cdots \gamma_{\sigma(j_{p})}$$

• write
$$\beta = \beta' \circ \sigma$$
, with $\epsilon(\beta) = \epsilon(\beta')\epsilon(\sigma)$

$$T_{\alpha} = \sum_{1 \leq j_1 < \dots < j_p \leq m} \sum_{\beta' \in S_p} \epsilon(\beta') \sum_{\sigma \in S_p} \epsilon(\sigma) \delta_{\sigma(j_1)}(\boldsymbol{a}_{\alpha}^{\beta'(\sigma(1))}) \cdots \delta_{\sigma(j_p)}(\boldsymbol{a}_{\alpha}^{\beta'(\sigma(p))}) \times \gamma_{\sigma(j_1)} \cdots \gamma_{\sigma(j_p)}$$

use commutativity of A to rearrange the $\delta_{\sigma(j_k)}(a_{\alpha}^{\beta'(\sigma(k))})$:

Operators

Algebra

Openness

Tentative charts

000000000

Reconstruction

$$T_{\alpha} = \sum_{1 \le j_1 < \dots < j_p \le m} \sum_{\beta' \in S_p} \epsilon(\beta') \delta_{j_1}(a_{\alpha}^{\beta'(1)}) \cdots \delta_{j_p}(a_{\alpha}^{\beta'(p)})$$
$$\times \sum_{\sigma \in S_p} \epsilon(\sigma) \gamma_{\sigma(j_1)} \cdots \gamma_{\sigma(j_p)}$$
$$= \sum_{1 \le j_1 < \dots < j_p \le m} \det(\delta_j a_{\alpha}^k) \sum_{\sigma \in S_p} \epsilon(\sigma) \gamma_{\sigma(j_1)} \cdots \gamma_{\sigma(j_p)}$$

0000000 000 00000

Recall the conditional expectation values $E_{\mathcal{A}}$: End_{\mathcal{A}}(\mathcal{H}_{∞}) $\rightarrow \mathcal{A}$ from the projections $p_j \in \mathcal{A}$ of Tr(e) = $\sum_{i=1}^{n} jp_i$ by

$$E_{\mathcal{A}}(T) := \sum_{k=1}^{n} \frac{p_k}{k} \sum_{j=1}^{n} T_{jj}, \qquad T = (T_{ij}) \in eM_n(\mathcal{A})e$$

Gelfand-Naimark

Serre-Swan

Spectral triples

Gelfand-Naimark	Serre-Swan	Spectral triples	Operators	HH	Algebra	Openness	Tentative charts	Reconstruction
							000000000	

Tentative charts

Definition

$$\mathbf{D} \ \rho_{\alpha} := \mathbf{i}^{\frac{\mathbf{p}(\mathbf{p}+1)}{2}} \mathbf{E}_{\mathcal{A}}(\gamma \mathbf{T}_{\alpha})$$

(no γ for p odd, by construction $\rho_{\alpha} = \rho_{\alpha}^* \in \mathcal{A}$)

$$2 \quad U_{\alpha} := \{\chi \in X : \rho_{\alpha}(\chi) \neq 0\} \subset X$$

Proposition

- The U_{α} form an open cover of X.
- Suppose all derivation of the form $\delta_j(a) = i(\eta_j | [D, a]\eta_j)$, for $\eta_j \in \mathcal{H}_{\infty}$, satisfy the Main Assumption. Let $\chi \in U_{\alpha}$. Then:
 - i) There exist p derivations $\delta_1, \ldots, \delta_p$ with det $(\chi(\delta_j a_{\alpha}^k)) \neq 0$
- ii) The map $\mathbf{s}_{\alpha} : U_{\alpha} \to \mathbb{R}^{p}$ is continuous and open.

Remarks on the main assumption

Spectral triples

By investigation of maps $t \mapsto \gamma_t(a) = e^{it|D|} a e^{-it|D|}$ one proves

0000000 000 00000

Algebra

Openness

Tentative charts

000000000

Reconstruction

Operators

Theorem

Serre-Swan

Gelfand-Naimark

For any $h = h^* \in A$ the commutator [D, h] vanishes where h reaches its maximum.

More precisely, for any sequence $b_n \in A$, with $||b_n|| \le 1$ and support tending to $\{\chi\}$, where χ is a character such that $|\chi(h)||$ is maximal, one has $||[D, h]b_n|| \to 0$.

It follows:

Proposition

The derivations $\pm \delta_j$, with $\delta_j(a) = i(\eta_j | [D, a] \eta_j)$, are dissipative, i.e.

$$\|oldsymbol{a}+\lambda\delta_j(oldsymbol{a})\|\geq\|oldsymbol{a}\|\quadoralloldsymbol{a}\in\mathcal{A}\;,\;\lambda\in\mathbb{R}$$

Gelfand-Naimark	Serre-Swan	Spectral triples	Operators	ΗH	Algebra	Openness	Tentative charts	Reconstruction
							000000000	

As a byproduct, one obtains

Proposition

For $h = h^* \in A$ one has, with norm convergence,

$$\lim_{ au o \infty} rac{{f e}^{{
m i} au {m h}} |{m D}| {m e}^{-{
m i} au {m h}}}{ au} \xi = |[{m D},{m h}]| \xi \;, \qquad \xi \in {
m dom}({m D})$$

- this implies [|D, h|, [D, a]] = 0 for all $a, h \in A$ with $h = h^*$
- [*D*, *a*][*D*, *b*] + [*D*, *b*][*D*, *a*] commute with *A* and [*D*, *A*]
- under the strong regularity assumption, the Clifford algebra is recovered: [D, a][D, b] + [D, b][D, a] ∈ A

```
Using dissipativity of \delta_0, Sobolev estimates and the Hille-Yosida-Lumer-Phillips Theorem, one shows that U(t) = e^{t\delta_0} is a one-parameter group of isometries of the C^*-algebra A.
```

It is further shown that this group preserves smoothness

Local bound for multiplicity

Spectral triples

Serre-Swan

The continuous open map $s_{\alpha} : U_{\alpha} \to \mathbb{R}^{p}$ is not injective. Show that at most finitely many $\chi_{i} \in U_{\alpha}$ map to the same point:

0000000 000 00000

Algebra

Openness

Tentative charts

Reconstruction

•••••

• the measure $F_t^*(\mu)$ is strongly equivalent to μ

Operators

- $s_{\alpha}(\mu)$ is locally equivalent to the Lebesgue measure on \mathbb{R}^{p}
- $\#\{s_{\alpha}^{-1}(y)\} \leq \Sigma(y)$ (Lebesgue almost everywhere), where $\Sigma(y)$ is joint spectral multiplicity of action of a_{α}^{k} on \mathcal{H}
- By results of Voiculescu, this is controlled by a norm on $\mathcal{L}^{(p,1)}$

Theorem

Gelfand-Naimark

Let $V \subset U_{\alpha}$ be an open set with $\overline{V} \subset U_{\alpha}$. Then there exists $m < \infty$ such that

$$\#\{s_{\alpha}^{-1}(y) \cap V\} < m \quad \forall y \in s_{\alpha}(V) \ .$$

Gelfand-Naimark	Serre-Swan	Spectral triples	Operators	ΗH	Algebra	Openness	Tentative charts	Reconstruction
								000000000

Proposition

Let $V \subset U_{\alpha}$ be an open set with $\overline{V} \subset U_{\alpha}$.

There exists a dense open subset $Y \subset s_{\alpha}(V)$ such that every point of $s_{\alpha}^{-1}(Y) \cap V$ has a neighbourhood *N* in *X* such that the restriction of s_{α} to *N* is an homeomorphism with an open set of \mathbb{R}^{p} .

Stratety:

•
$$m_1 := \sup_{y \in s_\alpha(V)} \#\{s_\alpha^{-1}(y) \cap V\}$$
, with $0 < m_1 < \infty$

•
$$Y_1 := \{ y \in s_{\alpha}(V) : \#\{s_{\alpha}^{-1}(y) \cap V\} = m_1$$

- Y₁ is open
- $\exists V_1, \ldots, V_{m_1} \subset V$, with V_i open and mutually disjoint, with $s_{\alpha}(V_i) = Y_1$

Reconstruction Theorem

Lemma

For every point $\chi \in X$ there exist

- *p* real elements $x^{\mu} \in \mathcal{A}$,
- a smooth family $\tau_t \in Aut(\mathcal{A}), t \in \mathbb{R}^p, \tau_0 = id$,

such that

- The x^μ give a homeomorphism of a neighbourhood of χ with an open set in ℝ^ρ.
- 2 The map t → h(t) = τ^{*}_t χ is a homeomorphism of a neighbourhood W of 0 in ℝ^p with a neighbourhood of χ.
- **③** The map $\psi = \chi \circ h : W \to \mathbb{R}^p$ is a local diffeomorphism.

Gelfand-Naimark	Serre-Swan	Spectral triples	Operators	HH	Algebra	Openness	Tentative charts	Reconstruction
								0000000000

Proposition

The algebra \mathcal{A} is locally the algebra of restrictions of smooth functions on \mathbb{R}^{p} to a bounded open set of \mathbb{R}^{p} .

(\Rightarrow) C^{∞} -functional calculus

Given selfadjoint $x_1, \ldots, x_p \in A$ and smooth function $f : \mathbb{R}^p \to \mathbb{C}$ defined on a neighbourhood of the joint spectrum of the x_j . Then

$$f_{(\boldsymbol{x}_1,\ldots,\boldsymbol{x}_p)} := \frac{1}{(2\pi)^p} \int_{\mathbb{R}^p} d(t_1,\ldots,t_p) \ \hat{f}(t_1,\ldots,t_p) \exp\left(\mathrm{i} \sum_{j=1}^p t_j \boldsymbol{x}_j\right) \in \mathcal{A}$$

(\Leftarrow) use the diffeomorphisms $\phi_{\kappa} : \mathbb{R}^{p} \supset W \rightarrow Y \subset \mathbb{R}^{p}$

Theorem

Let $(\mathcal{A}, \mathcal{H}, D)$ be a strongly regular spectral triple fulfilling the five conditions, with c antisymmetric. Then there exists an oriented smooth compact manifold X such that $\mathcal{A} = C^{\infty}(X)$.

- change of charts $x_2 : V_2 \to \mathbb{R}^p$ to $x_1 : V_1 \to \mathbb{R}^p$, with $\chi \in V_1 \cap V_2$
- By previous proposition, there exist *p* smooth functions $f^{\mu} \in C^{\infty}(x_2(V_1 \cap V_2))$ with $x_1^{\mu}(\chi) = (f^{\mu} \circ x_2)(\chi)$.

• Let
$$y = x_2(\chi) \in x_2(V_1 \cap V_2)$$
. Then

$$x_1^{\mu} \circ x_2^{-1}(y) = x_1^{\mu}(\chi) = f^{\mu}(y)$$

i.e. $x_1 \circ x_2^{-1}$ is smooth

• Hochschild cycle c gives a nowhere vanishing section of $\Lambda^{p}(T^{*}X)$

Gelfand-Naimark	Serre-Swan	Spectral triples	Operators	HH	Algebra	Openness	Tentative charts	Reconstruction
								0000000000

The converse

Theorem

An involutive algebra \mathcal{A} is the algebra of smooth functions on an oriented smooth compact manifold if and only if it admits a faithful representation in a pair (\mathcal{H}, D) fulfilling the five conditions of a spectral triple with the Hochschild cycle antisymmetric and the strong regularity.

- $\mathcal{H} = L^2(X, \Lambda^*_{\mathbb{C}})$
- D = d + d* for codifferential d* with respect to any metric on X

•
$$[D, f]\xi = (\partial_{\mu}f)\gamma^{\mu}$$
 with $\gamma^{\mu}\xi = e_{\mu} \wedge \xi - i_{e_{\mu}}\xi$

•
$$\gamma = i^{-\frac{p(p+1)}{2}} e^1 \wedge \cdots \wedge e^p$$

regularity:

use
$$(D^2 + 1)^{\frac{1}{2}} =: \langle D \rangle = \frac{1}{\pi} \int_0^\infty \frac{d\lambda}{\sqrt{\lambda}} \frac{\langle D \rangle^2}{\lambda + \langle D \rangle^2}$$
 to obtain

$$\begin{split} \tilde{\delta}^m T &= \frac{1}{\pi^m} \int_0^\infty \Big(\prod_{i=1}^m \frac{1}{\langle D \rangle^2 + \lambda_i} \Big) ((\operatorname{ad} D^2)^m T) \Big(\prod_{j=1}^m \frac{d\lambda_j \sqrt{\lambda_j}}{\langle D \rangle^2 + \lambda_j} \Big) \\ &= \sum_{k=0}^m (-1)^k \binom{m}{k} \frac{1}{\pi^m} \int_0^\infty \Big(\prod_{i=1}^k \frac{1}{\langle D \rangle^2 + \lambda_i} \Big) ((\operatorname{ad} D^2)^{m+k} T) \langle D \rangle^{-m} \\ &\times \prod_{j=1}^m \frac{d\lambda_j \sqrt{\lambda_j} \langle D \rangle}{(\langle D \rangle^2 + \lambda_j)^2} \end{split}$$

- D^2 is scalar elliptic operator with principal symbol $g_{\mu
 u}$
- Let P be a ΨDO of order q, then the principal symbol of [D², P] of order q + 2 vanishes

Gelfand-Naimark	Serre-Swan	Spectral triples	Operators	ΗН	Algebra	Openness	Tentative charts	Reconstruction
								00000000000

Spin^c-manifolds

Definition

Let (X, g) be a smooth compact Riemannian manifold and A = C(X). Let $B = \Gamma(C\ell(X))$ be the algebra of continuous sections of the Clifford bundle over X, generated by T^*X and g^{-1} .

A Clifford module over X is a finitely generated projective right A-module \mathcal{E} with hermitian structure $(|) : \mathcal{E} \times \mathcal{E} \to A$, together with a homomorphism $c : B \to \operatorname{End}_A(\mathcal{E})$, such that $(\xi | c(\lambda)\eta) = (c(\lambda^*)\xi | \eta)$ for $\lambda \in B$ and $\xi, \eta \in \mathcal{E}$. (This makes \mathcal{E} a *B*-*A*-bimodule.)

X is called a spin^{*c*}-manifold if there is a *B*-*A*-bimodule \mathcal{E} with $\operatorname{End}_{A}(\mathcal{E}) \simeq B$, and each isomorphism class of such *B*-*A*-bimodules is called a spin^{*c*}-structure.



Theorem

Let $(\mathcal{A}, \mathcal{H}, D)$ be a spectral triple, with \mathcal{A} commutative and the cycle *c* antisymmetric. Assume that the multiplicity of the action of \mathcal{A}'' in \mathcal{H} is $2^{\left[\frac{p}{2}\right]}$. Then there exists a smooth oriented compact spin^c-manifold X such that $A = C^{\infty}(X)$.

- Starting point: [([D, a][D, b] + [D, b][D, a]), [D, c]] = 0 for all a, b, c ∈ A
- for dimensional reasons: $\operatorname{End}_{\mathcal{H}} = \Gamma^{\infty}(C\ell_Q(X))$
- $c: (da)(\chi) \rightarrow i[D, a](\chi)$ isomorphism
- [D, a][D, b] + [D, b][D, a] =: -2g^{-1}(da, db) metric
- $\|da\| = \sup_{\chi \in X} \|(\operatorname{grad} a)(\chi)\|$ leads to distance formula

 $\operatorname{dist}_{\boldsymbol{g}}(\boldsymbol{\chi},\boldsymbol{\kappa}) = \sup\{|\boldsymbol{a}(\boldsymbol{\chi}) - \boldsymbol{a}(\boldsymbol{\kappa})| \ : \ \boldsymbol{a} \in \mathcal{A} \ , \ \|[\mathcal{D},\boldsymbol{a}]\| \leq 1\}$



- We have sketched the proof of a recent Theorem of A. Connes which establishes a 1:1 correspondence between commutative spectral triples and smooth compact oriented manifolds.
- The purpose was to present parts of the enormous diversity of methods used in Noncommutative Geometry.
- The reconstruction theorem provides strong motivation for noncommutative spectral triples as possible candidates for new forms of geometry in the early universe.