

Sheaf theory for stacks in manifolds and twisted cohomology for S^1 -gerbes

Ulrich Bunke, Thomas Schick and Markus Spitzweck *

January 17, 2007

Contents

1	Introduction	1
1.1	About the motivation	1
1.2	Statement of the main result	2
1.3	Twisted cohomology and gerbes	3
1.4	Sheaf theory for smooth stacks	9
2	Sheaf theory for smooth stacks	11
2.1	Over sites	11
2.2	The site of a smooth stack	14
2.3	Presheaves of abelian groups and derived functors	22
2.4	Simplicial models	24
2.5	Comparison of big and small sites	29
3	The de Rham complex	32
3.1	The de Rham complex is a flabby resolution	32
3.2	Calculation for $U(1)$ -gerbes	34

1 Introduction

1.1 About the motivation

1.1.1 Given a closed three form $\lambda \in \Omega^3(X)$ on a smooth manifold X , the usual definition of twisted de Rham cohomology is as the cohomology of the two-periodic complex $(\Omega_{per}^\bullet(X), d_\lambda)$, where

$$\Omega_{per}^\bullet(X) := \bigoplus_{n \in \mathbb{Z}} \Omega^{\bullet+2n}(X), \quad \text{and} \quad d_\lambda := d_{dR} + \lambda$$

is the sum of the de Rham differential and the multiplication operator by the form λ .

1.1.2 Twisted de Rham cohomology is in particular interesting as a target of the Chern character from twisted K -theory. In this case $[\lambda] \in H^3(X; \mathbb{R})$ is the real image of an integral class $\lambda_{\mathbb{Z}}(P) \in H^3(X; \mathbb{Z})$ which classifies a principal bundle $P \rightarrow X$ with structure group PU , the projective unitary group of a complex infinite-dimensional separable Hilbert space. The twisted K -theory depends functorially on P in a non-trivial manner.

*Mathematisches Institut, Universität Göttingen, Bunsenstr. 3-5, 37073 Göttingen, GERMANY, bunke@uni-math.gwdg.de, schick@uni-math.gwdg.de, spitz@uni-mah.gwdg.de

The twisted cohomology as defined above depends on the cohomology class $[\lambda]$ up to (in general) non-canonical isomorphism. The draw-back of this definition of twisted cohomology above is that it is not functorial in the twist $P \rightarrow X$ of K -theory since there no canonical choice of a three-form λ representing the image of $\lambda_{\mathbb{Z}}(P)$ in real cohomology.

1.1.3 The main goal of the present note is to propose an alternative functorial definition of the twisted cohomology as the real cohomology of a stack G_P which is canonically associated to the PU -bundle $P \rightarrow X$. The stack G_P is the stack of U -reductions of $P \rightarrow X$, where U is the unitary group of the Hilbert space and $U \rightarrow PU$ is the canonical projection map. It is also called the lifting gerbe of P .

In order to define the cohomology of a stack like G_P we develop a sheaf theory set-up for stacks in smooth manifolds. Our main result Theorem 1.1 is the key step in the verification that the cohomology according to the new sheaf-theoretic definition is essentially isomorphic (non-canonically) to the twisted cohomology as defined above.

We have chosen to work with stacks in smooth manifolds since we are heading towards a comparison with de Rham cohomology. A parallel theory can be set up in the topological context. Together with applications to T -duality and delocalized cohomology it will be discussed in detail in the subsequent papers [8], [6] and [7].

1.1.4 The twists for our new cohomology theory are smooth gerbes $G \rightarrow X$ with band $U(1)$. The lifting gerbe $G_P \rightarrow X$ of a PU -bundle mentioned above is an example. Advantages of our new definition are:

- (1) The twisted cohomology depends functorially on the twist.
- (2) One can define twisted cohomology with coefficients in an arbitrary abelian group.
- (3) The definition can easily be generalized to the topological context.

1.1.5 In Subsection 1.2 we give a complete technical statement of our main result written for a reader familiar with the language of stacks, sites, and sheaf theory. The third part of the introduction, Subsection 1.3, is devoted to a detailed motivation with references to the literature and a less technical introduction of the language and the description of the result. Finally, Subsection 1.4 is an introduction to the technical sheaf theoretic part of the present paper.

1.2 Statement of the main result

1.2.1 We consider a stack G on the category of smooth manifolds equipped with the usual topology of open coverings. To G we associate a site \mathbf{G} as a subcategory of manifolds over G . The objects of this site are representable smooth maps $U \rightarrow G$ from smooth manifolds to G . A covering $(U_i \rightarrow U)_{i \in I}$ is a collection of morphisms which are submersions and such that $\sqcup_{i \in I} U_i \rightarrow U$ is surjective (see 2.2.3 for a precise definition).

1.2.2 To the site \mathbf{G} we associate the categories of presheaves $\text{Pr } \mathbf{G}$ and sheaves $\text{Sh } \mathbf{G}$ of sets as well as the lower bounded derived categories $D^+(\text{Pr}_{\text{Ab}} \mathbf{G})$ and $D^+(\text{Sh}_{\text{Ab}} \mathbf{G})$ of the abelian categories $\text{Pr}_{\text{Ab}} \mathbf{G}$ and $\text{Sh}_{\text{Ab}} \mathbf{G}$ of presheaves and sheaves of abelian groups.

1.2.3 Let $i : \text{Sh } \mathbf{G} \rightarrow \text{Pr } \mathbf{G}$ be the natural inclusion, and let $i^\sharp : \text{Pr } \mathbf{G} \rightarrow \text{Sh } \mathbf{G}$ be its left adjoint, the sheafification functor. As a right adjoint the functor i is left exact and admits a right derived functor $Ri : D^+(\text{Sh}_{\text{Ab}} \mathbf{G}) \rightarrow D^+(\text{Pr}_{\text{Ab}} \mathbf{G})$.

1.2.4 If $G \rightarrow X$ is a morphism of stacks, then we define a functor $f_* : \text{Pr } \mathbf{G} \rightarrow \text{Pr } \mathbf{X}$. Note that if f is not representable, then this map is not associated to a map of sites. If $F \in \text{Pr } \mathbf{G}$ and $(U \rightarrow X) \in \mathbf{X}$, then we set (see 2.4)

$$f_*(U) := \lim F(V) ,$$

where the limit is taken over the category of diagrams

$$\begin{array}{ccc} V & \longrightarrow & G \\ \downarrow & \nearrow & \downarrow f \\ U & \longrightarrow & X \end{array} .$$

It turns out that f_* admits a left adjoint. Therefore it is left exact and admits a right derived functor $Rf_* : D^+(\mathrm{Pr}_{\mathrm{Ab}} \mathbf{G}) \rightarrow D^+(\mathrm{Pr}_{\mathrm{Ab}} \mathbf{X})$.

1.2.5 Let $f : G \rightarrow X$ be a smooth gerbe with band S^1 over the smooth manifold X . We consider the sheafification $i^\sharp \mathbb{R}_{\mathbf{G}}$ of the constant presheaf $\mathbb{R}_{\mathbf{G}}$ on \mathbf{G} with value \mathbb{R} . Our main result describes

$$i^\sharp \circ Rf_* \circ Ri(i^\sharp \mathbb{R}_{\mathbf{G}}) \in D^+(\mathrm{Sh}_{\mathrm{Ab}} \mathbf{X})$$

in terms of a deformation of the de Rham complex.

The gerbe $f : G \rightarrow X$ is classified by a Dixmier-Douady class $\lambda_{\mathbb{Z}} \in H^3(X; \mathbb{Z})$. Let $\lambda \in \Omega^3(X)$ be a closed form such that $[\lambda] \in H^3(X; \mathbb{R})$ represents the image of $\lambda_{\mathbb{Z}}$ under $H^3(X; \mathbb{Z}) \rightarrow H^3(X; \mathbb{R})$.

For a manifold X the objects (U, p) of the site \mathbf{X} are submersions $p : U \rightarrow X$ from smooth manifolds U to X . This differs from the usual convention, where the site is the category of open subsets of X .

We form the complex of presheaves $(U, p) \mapsto \Omega[[z]]_\lambda(U, p)$ on \mathbf{X} , which associates to $(U, p) \in \mathbf{X}$ the complex of formal power series of smooth real differential forms on U with differential

$$d_\lambda := d_{dR} + T\lambda,$$

where z is a formal variable of degree 2, $T := \frac{d}{dz}$, d_{dR} is the deRham differential, and λ stands for multiplication by $p^*\lambda$. It turns out that this is actually a complex of sheaves (see Lemma 3.1).

1.2.6 The main result of the present paper is the following theorem.

Theorem 1.1 *In $D^+(\mathrm{Sh}_{\mathrm{Ab}} \mathbf{X})$ we have an isomorphism $i^\sharp \circ Rf_* \circ Ri(i^\sharp \mathbb{R}_{\mathbf{G}}) \cong \Omega[[z]]_\lambda$.*

1.2.7 The projection map $f : G \rightarrow X$ of a gerbe is not representable so that $f_* : \mathrm{Pr} \mathbf{G} \rightarrow \mathrm{Pr} \mathbf{X}$ does not come from an associated map of sites. Therefore, in order to define Rf_* and to verify the theorem we have to develop some standard elements of sheaf theory for stacks in smooth manifolds. This is the contents of Section 2 (see 1.4 for an introduction). In Section 3 we verify Theorem 1.1.

1.3 Twisted cohomology and gerbes

1.3.1 A closed three-form $\lambda \in \Omega^3(X)$ on a smooth manifold X can be used to perturb the de Rham differential

$$d_{dR} \rightsquigarrow d_{dR} + \lambda =: d_\lambda.$$

The cohomology of the two-periodic complex

$$\dots \xrightarrow{d_\lambda} \Omega^{\mathrm{even}}(X) \xrightarrow{d_\lambda} \Omega^{\mathrm{odd}}(X) \xrightarrow{d_\lambda} \Omega^{\mathrm{even}}(X) \xrightarrow{d_\lambda} \dots$$

is called the λ -twisted cohomology of X and often denoted by $H^*(X; \lambda)$. This ad-hoc definition appears in various places in the recent mathematical literature (let us mention just [1], [3], [16], [4]) and in the physics literature. A closely related and essentially equivalent definition [15] uses the complex $(\Omega(X)((u)), d_{dR} - u\lambda)$, where u is a formal variable of degree -2 , and $((u))$ stands for formal Laurent series.

1.3.2 It is known that the isomorphism class of the λ -twisted cohomology group only depends on the cohomology class $[\lambda] \in H^3(X; \mathbb{R})$. If $f : Y \rightarrow X$ is a smooth map, then we have a functorial map $f^* : H^*(X; \lambda) \rightarrow H^*(Y; f^*\lambda)$ which essentially only depends on the homotopy class of f . Furthermore, λ -twisted cohomology has a Mayer-Vietoris sequence and is a module over $H^*(X; \mathbb{R})$. It now appears as a natural question to understand λ -twisted cohomology as a concept of algebraic topology.

1.3.3 One attempt is the approach of [9] in which the complex of smooth differential forms is replaced by similar objects in algebraic topology.

The proposal of [1] to use the singular de Rham complex goes into the same direction. Observe that we can use the filtration of $\Omega^{\mathrm{ev}}(X)$ and $\Omega^{\mathrm{odd}}(X)$ by degree in order to construct a spectral sequence converging to $H^*(X; \lambda)$. Its E_2 -page involves $H^*(X; \mathbb{R})$ (as $\mathbb{Z}/2\mathbb{Z}$ -graded vector spaces). The next possibly non-trivial differential of this spectral sequence is the multiplication by the class $[\lambda]$. In [1] the higher differentials of this spectral sequence are identified as Massey products.

1.3.4 A natural homotopy theoretic framework for twisted cohomology theories would be some version of parametrized stable homotopy theory as developed e.g. in [17]. In such a theory a twist of a generalized cohomology theory (represented by a spectrum E) is a parametrized spectrum \mathcal{E} over X with typical fibre equivalent to E (think of a bundle of spectra). The twisted cohomology groups $H^*(X; \mathcal{E})$ are then given by the homotopy groups of the spectrum of sections of \mathcal{E} . In order to interpret λ -twisted cohomology in this manner one would have to relate three-forms on X with parametrized versions of the Eilenberg-MacLane spectrum $H\mathbb{R}$.

Let us mention that alternatively to [17] other reasonable versions of a stable homotopy theory over X could be based on presheaves of spectra over X or $\Omega(X)$ -equivariant spectra, where $\Omega(X)$ denotes the based loops of X .

1.3.5 One motivation for introducing λ -twisted cohomology is based on the fact that it can be used as a target of the Chern character from twisted K -theory. It is known that $H^3(X; \mathbb{Z})$ classifies a certain subset of isomorphism classes of parametrized spectra \mathcal{K} with fibre equivalent to the complex K -theory spectrum K . This follows from the splitting $BGL_1(K) \cong K(\mathbb{Z}, 3) \wedge T$. Here $GL_1(K)$ denotes the grouplike monoid of units of the K -theory spectrum, $K(\mathbb{Z}, 3)$ denotes an Eilenberg-MacLane space, and T is an auxiliary space. We refer to [18] for more details. Chern characters are constructed in [3], [1], [16], [15]. Note that in these works twisted K -theory is not defined in homotopy theoretic terms but using sections in bundles of Fredholm operators, bundle gerbe modules or K -theory of C^* -algebras. If $\lambda_{\mathbb{Z}} \in H^3(X; \mathbb{Z})$ classifies the parametrized K -theory spectrum \mathcal{K} , then the Chern character has values in $H^*(X; \lambda)$, where $[\lambda]$ is the image of $\lambda_{\mathbb{Z}}$ under the map $H^3(X; \mathbb{Z}) \rightarrow H^3(X; \mathbb{R})$. Such a definition can not be natural since in general \mathcal{K} has non-trivial automorphisms which are not reflected by $H^3(X; \lambda)$.

A completely natural definition of a Chern character with values even in a twisted rational cohomology could be induced from the canonical rationalization map $\mathcal{K} \rightarrow \mathcal{K}_{\mathbb{Q}}$ if we like to define twisted rational cohomology using $\mathcal{K}_{\mathbb{Q}}$.

1.3.6 Above we have seen that $H^3(X; \mathbb{Z})$ classifies a subset of the isomorphism classes of parametrized K -theory spectra over X . This can in fact be seen directly. Let U be the unitary group of a separable infinite dimensional complex Hilbert space. Equipped with the topology induced by the operator norm it is a topological group. By Kuipers theorem it is contractible so that the projective unitary group $PU := U/U(1)$ has the homotopy type of $BU(1) \cong K(\mathbb{Z}, 2)$. Taking the classifying space once more we have $BPU \cong K(\mathbb{Z}, 3)$. This shows that $H^3(X; \mathbb{Z})$ classifies isomorphism classes of PU -principal bundles over X . One can now manufacture a PU -equivariant version of a K -theory spectrum K (see e.g. [12]). If $P \rightarrow X$ is a PU -principal bundle, then one can define the bundle of spectra $\mathcal{K} := P \times_{PU} K$ over X . Alternatively one could construct twisted K -theory starting from a bundle of projective Hilbert spaces as in [2]. As a result of this discussion one should consider PU -principal bundles as more primary objects.

1.3.7 The theory of bundle gerbes initiated in [20] and continued in [21] aims at a categorification of $H^3(X; \mathbb{Z})$ in a similar manner as $U(1)$ -principal bundles categorify $H^2(X; \mathbb{Z})$. The PU -principal bundles considered above are particularly nice examples of bundle gerbes. Other examples of bundle gerbes are introduced in [11]. In order to simplify we forget the smooth structure of X for the moment and work in the category of topological spaces.

Let us represent X as a moduli space of a groupoid $A^1 \rightrightarrows A^0 \rightarrow X$ in topological spaces, i.e. we represent X as the quotient of the space of objects A^0 by the equivalence relation A^1 . In addition we shall assume that the range and source maps have local sections. Then a bundle gerbe is the same as a central $U(1)$ -extension

$$\begin{array}{ccc}
 U(1) & & \\
 \downarrow & & \\
 \tilde{A}^1 & \rightrightarrows & A^0 \\
 \downarrow & & \parallel \\
 A^1 & \rightrightarrows & A^0 \longrightarrow X
 \end{array}$$

of topological groupoids.

In order to relate the PU -principal bundle $P \rightarrow X$ with a bundle gerbe we represent X as the moduli space of the action groupoid $P \times PU \rightrightarrows P \rightarrow X$. The central $U(1)$ -central extension of this groupoid is given by $P \times U \rightrightarrows P$.

1.3.8 The picture of a gerbe in [11] is obtained by choosing an open covering $(U_i)_{i \in I}$ of X and forming the representation

$$\bigsqcup_{i,j} U_i \cap U_j \rightrightarrows \bigsqcup_{i \in I} U_i \rightarrow X .$$

The data of a $U(1)$ -central extension of this groupoid is equivalent to transition line bundle data and trivializations over triple intersection considered in [11].

One can build a two-category of topological groupoids by inverting Morita equivalence such that equivalence classes of $U(1)$ -central extensions of groupoids representing X are indeed classified by $H^3(X; \mathbb{Z})$ (see e.g. [25]).

1.3.9 A more natural view on this category of groupoids is through stacks on topological spaces \mathbf{Top} . We consider \mathbf{Top} as a Grothendieck site where covering families are given by coverings by families of open subspaces.

Note that groupoids form a two-category. A stack G on \mathbf{Top} can be viewed as an object which associates to each space $U \in \mathbf{Top}$ a groupoid $G(U)$, to a morphism $U' \rightarrow U$ a homomorphism of groupoids $G(U) \rightarrow G(U')$, to a chain of composeable morphisms

$$\begin{array}{ccc} U'' & \longrightarrow & U \\ & \searrow & \nearrow \\ & U' & \end{array}$$

a two-isomorphism

$$\begin{array}{ccc} G(U) & \longrightarrow & G(U'') \\ & \searrow & \nearrow \\ & G(U') & \end{array}$$

satisfying a natural associativity relation, and such that G satisfies descent conditions for the covering families of U . Precise definitions can be found e.g. in [22], [10], [5]. A space $V \in \mathbf{Top}$ can be viewed as a stack by the Yoneda embedding such that $V(U) = \text{Hom}_{\mathbf{Top}}(U, V)$ (where we consider sets as groupoids with only identity morphisms).

1.3.10 As an illustration we explain a canonical construction which associates to a PU -principal bundle $P \rightarrow X$ over a space X a stack G_P together with a map $G_P \rightarrow X$. It will be called the lifting gerbe of P . Observe that U acts on P via the canonical homomorphisms $U \rightarrow PU$. For a space $T \in \mathbf{Top}$ the objects of the groupoid $G_P(T)$ are the diagrams

$$\begin{array}{ccc} Q & \longrightarrow & P \\ \downarrow & & \downarrow \\ T & \longrightarrow & X \end{array} ,$$

where $Q \rightarrow T$ is a U -principal bundle, and $Q \rightarrow P$ is U -equivariant.

A morphism between two such objects

$$\begin{array}{ccc} Q & \longrightarrow & P \\ \downarrow & & \downarrow \\ T & \longrightarrow & X \end{array} , \quad \begin{array}{ccc} Q' & \longrightarrow & P \\ \downarrow & & \downarrow \\ T & \longrightarrow & X \end{array} ,$$

is an isomorphism of U -principal bundles $Q \rightarrow Q'$ over T which is compatible with the maps to P .

Finally, for a map $T' \rightarrow T$ the functor $G_P(f) : G_P(T) \rightarrow G_P(T')$ maps the object

$$\begin{array}{ccc} Q & \longrightarrow & P \in G_P(T) \\ \downarrow & & \downarrow \\ T & \longrightarrow & X \end{array}$$

to the induced diagram

$$\begin{array}{ccc} T' \times_T Q & \longrightarrow & P \in G_P(T') , \\ \downarrow & & \downarrow \\ T' & \longrightarrow & X \end{array}$$

and a morphism $Q \rightarrow Q'$ to the induced morphism $T' \times_T Q \rightarrow T' \times_T Q'$. We leave it as an exercise to check that this presheaf of groupoids is a stack.

The morphism $G_P \rightarrow X$ maps the object

$$\begin{array}{ccc} Q & \longrightarrow & P \in G_P(T) \\ \downarrow & & \downarrow \\ T & \longrightarrow & X \end{array}$$

to the underlying map $T \rightarrow X$ which is considered as an element of $X(T)$.

1.3.11 A diagram of PU -principal bundles

$$\begin{array}{ccc} P & \longrightarrow & P' \\ \downarrow & & \downarrow \\ X & \longrightarrow & X' \end{array}$$

functorially induces a diagram of stacks

$$\begin{array}{ccc} G_P & \longrightarrow & G_{P'} \\ \downarrow & & \downarrow \\ X & \longrightarrow & X' \end{array}$$

in the obvious way.

1.3.12 A topological groupoid $A : A^1 \rightrightarrows A^0$ represents a stack $[A^1/A^0]$ on topological spaces. It associates to each space U the groupoid $[A^0/A^1](U) = \text{Hom}(U, [A^0/A^1])$ of A -principal bundles on X and isomorphisms (see [10]). A morphism of groupoids gives rise via an associated bundle construction to a map of stacks. In this way we can embed the two-category of topological groupoids (with Morita equivalence inverted) mentioned at the end of 1.3.7 as a full subcategory of stacks on Top . The image of this embedding consists of topological stacks G , i.e. stacks which admit an atlas $A^0 \rightarrow G$.

An atlas is a surjective representable morphism $A^0 \rightarrow G$ admitting local sections, where A^0 is a space. Given an atlas of G we can construct a groupoid $A^1 \rightrightarrows A^0$. The morphism space of the groupoid is given by $A^1 := A^0 \times_G A^0$. We then have an equivalence of stacks $[A^0/A^1] \cong G$.

A map of stacks $G \rightarrow H$ is called representable if for any map $U \rightarrow H$ with U a space $U \times_H G$ is equivalent to a space. The representability condition on $A^0 \rightarrow G$ ensures that $A^1 := A^0 \times_G A^0$ is a space.

1.3.13 The lifting gerbe G_P of a PU -principal bundle 1.3.10 is a topological stack. In order to construct an atlas we choose a covering of X by open subsets on which P is trivial. Let A be the disjoint union of the elements of the covering, and $A \rightarrow X$ be the canonical map. By choosing local trivializations we

obtain the lift in the diagram

$$\begin{array}{ccc} & & P \\ & \nearrow s & \downarrow \\ A & \longrightarrow & X \end{array} .$$

We now consider the diagram

$$\begin{array}{ccc} A \times U & \xrightarrow{\phi} & P \in G_P(A) , \\ \downarrow & & \downarrow \\ A & \longrightarrow & X \end{array}$$

where $\phi(a, u) := s(a)\bar{u}$ and \bar{u} denotes the image of $u \in U$ under $U \rightarrow PU$. We consider this object as a morphism $A \rightarrow G_P$. We leave it as an exercise to verify that this map is an atlas.

1.3.14 A morphism of stacks $G \rightarrow X$ with X a space is a topological gerbe with band $U(1)$ if there exists an atlas $A \rightarrow X$, a lift

$$\begin{array}{ccc} & & G \\ & \nearrow & \downarrow \\ A & \longrightarrow & X \end{array} \tag{1.2}$$

to an atlas of G such that

$$\begin{array}{ccccc} & & U(1) & & \\ & & \downarrow & & \\ & & A \times_G A & \xrightarrow{\quad} & A \\ & & \downarrow & & \parallel \\ & & A \times_X A & \xrightarrow{\quad} & A \longrightarrow X \end{array}$$

is a $U(1)$ -extension of topological groupoids. In particular, the bundle gerbes considered in 1.3.7 give rise to topological gerbes with band $U(1)$. For equivalent definitions see [22], [10]. The definition of a gerbe in [5] is slightly more general since the existence of an atlas is not required.

1.3.15 The lifting gerbe $G_P \rightarrow X$ constructed in 1.3.10 is a topological gerbe with band $U(1)$. In fact, the construction 1.3.13 produces the lift (1.2).

1.3.16 In the definitions above the Grothendieck site \mathbf{Top} can be replaced by the Grothendieck site of smooth manifolds \mathbf{Mf}^∞ . In this site the covering families are again coverings by families of open submanifolds.

Stacks on \mathbf{Mf}^∞ are called stacks in smooth manifolds. If G is a stack in smooth manifolds, then an atlas $A \rightarrow G$ is a map of stacks which is representable and smooth, i.e. for any map $T \rightarrow G$ from a smooth manifold T to G the induced map $T \times_G A \rightarrow A$ is a submersion of manifolds. A stack in smooth manifolds which admits an atlas will then be called smooth.

1.3.17 Let $Y \rightarrow X$ be a map of manifolds. It is representable as a map between stacks in smooth manifolds if for any map $Z \rightarrow X$ the fibre product $Z \times_X Y$ exists as a manifold. Submersions between manifolds are representable maps.¹

1.3.18 We come to the conclusion that a basic object classified by $\lambda_{\mathbb{Z}} \in H^3(X; \mathbb{Z})$ is the equivalence class of a smooth gerbe $f : G \rightarrow X$ with band $U(1)$. Instead of going the way through some version of parametrized stable homotopy theory it now seems natural to define a real cohomology twisted by G directly using a suitable sheaf theory on stacks. A natural candidate would be something like $H^*(X; G) := H^*(G; \mathbb{R}) := H^*(G; i^\sharp \mathbb{R}_G)$, where $i^\sharp \mathbb{R}_G$ is the sheafification of the constant presheaf with value \mathbb{R} , and $H^*(\dots, i^\sharp \mathbb{R}_G)$ is defined using the derived global sections, or the derived p_* , where $p : G \rightarrow *$ is the

¹We do not know the converse, i.e. whether a representable map between manifolds is necessarily a submersion.

projection to a point. In fact, if G would be a manifold, then the sheaf theoretic $H^*(G, i^\sharp \mathbb{R}_G)$ would be isomorphic to the de Rham cohomology of the manifold G , and therefore to the topologist's $H^*(G; \mathbb{R})$. To proceed in the case of stacks we must clarify what we mean by a sheaf on G , and how we define p_* . The construction of $H^*(G; \mathbb{R})$ will be finalized in Definition 2.31. In order to define sheaves and presheaves on G we associate in 2.2 to G a Grothendieck site \mathbf{G} . The notions of presheaves and sheaves on a site are the standard ones.

1.3.19 If $f : G \rightarrow X$ is a morphism of stacks, then we are interested in functors f_*, f^* . Such operations are usually obtained from some induced morphisms of sites $f^\sharp : \mathbf{X} \rightarrow \mathbf{G}$. In fact, this works well for representable morphisms. But in the case of a gerbe $f : G \rightarrow X$ neither f nor $p : G \rightarrow *$ are representable. We will define f_* and p_* in an ad-hoc way. The same problem with a similar solution also occurs in algebro-geometric set-ups, see e.g. [14]. Because of this ad-hoc definitions we must redevelop some of the basic material of sheaf theory in order to check that the expected properties hold in the present set-up. For details we refer to the introduction 1.4 to sheaf theory part of the present paper.

1.3.20 After the development of elements of sheaf theory on smooth stacks we can define

$$H^*(X; G) := H^*(G; \mathbb{R}) := H^*(\mathbf{ev} \circ R p_* \circ Ri(i^\sharp \mathbb{R}_G)) ,$$

where $i : \mathbf{ShG} \rightarrow \mathbf{PrG}$ is the embedding of sheaves into presheaves, the sheafification functor $i^\sharp : \mathbf{PrG} \rightarrow \mathbf{ShG}$ is the left adjoint of i , and the exact functor

$$\mathbf{ev} : \mathbf{PrAbSite}(\ast) \rightarrow \mathbf{Ab}$$

evaluates a presheaf of abelian groups on the object $(\ast \rightarrow \ast) \in \mathbf{Site}(\ast)$. This last evaluation is necessary since our site is the big site of \ast consisting of all smooth manifolds. As the notation suggests we view this as the cohomology of X twisted by the gerbe G .

1.3.21 This definition is natural in G . If $u : G' \rightarrow G$ is a map of stacks, then by Lemma 2.16 we have a functorial map

$$u^* : H^*(G; \mathbb{R}) \rightarrow H^*(G'; \mathbb{R})$$

since there is a canonical isomorphism $u^* i^\sharp \mathbb{R}_G \cong i^\sharp \mathbb{R}_{G'}$. In particular, $H^*(X; G)$ carries the action of the automorphisms of the gerbe $G \rightarrow X$.

1.3.22 The natural question is now how the λ -twisted de Rham cohomology $H^*(X; \lambda)$ and $H^*(X; G)$ are related. The main step in this relation is provided by Theorem 1.1. Using this result in the isomorphism \cong and the projection $q : X \rightarrow \ast$ we can write

$$\begin{aligned} H^*(X; G) &= H^*(\mathbf{ev} \circ R p_* \circ Ri(i^\sharp \mathbb{R}_G)) \\ &\cong H^*(\mathbf{ev} \circ R(q \circ f)_* \circ Ri(i^\sharp \mathbb{R}_G)) \\ &\stackrel{(**)}{\cong} H^*(\mathbf{ev} \circ R q_* \circ R f_* \circ Ri(i^\sharp \mathbb{R}_G)) \\ &\stackrel{(*)}{\cong} H^*(\mathbf{ev} \circ R q_* \circ Ri \circ i^\sharp \circ R f_* \circ Ri(i^\sharp \mathbb{R}_G)) \\ &\stackrel{!}{\cong} H^*(\mathbf{ev} \circ R q_* \circ Ri(\Omega[[z]]_\lambda)) \\ &\stackrel{****)}{\cong} H^*(\mathbf{ev} \circ R(q_* \circ i)(\Omega[[z]]_\lambda)) \\ &\stackrel{(***)}{\cong} H^*(\mathbf{ev} \circ q_* \circ i(\Omega[[z]]_\lambda)) \\ &= H^*(\Omega[[z]]_\lambda(X)) . \end{aligned}$$

In order to justify the isomorphism $(*)$ we use Lemma 2.13 which says that f_* preserves sheaves. The isomorphism $(**)$ follows from Lemma 2.26 since f is smooth. For $(***)$ we use Lemma 2.27. Finally, $(****)$ follows from Lemma 2.30 and the fact that $\Omega[[z]]_\lambda$ is a complex of flabby sheaves (see 2.28).

Note that the isomorphism $\stackrel{!}{\cong}$ depends on additional choices.

1.3.23 It remains to relate the cohomology of the complex $(\Omega[[z]]_\lambda(X), d_\lambda)$ (see 1.2.5) with $H^*(X; \lambda)$. Let

$$\Omega[[z]]_\lambda(X)^p \subset \Omega[[z]]_\lambda(X)$$

be the subset of polynomials $\sum_{2n+k=p} z^n \omega^k$ with $\omega^k \in \Omega^k(X)$. Then we have $d_\lambda : \Omega[[z]]_\lambda(X)^p \rightarrow \Omega[[z]]_\lambda(X)^{p+1}$. For $p > 0$ we construct morphisms ψ_p such that the following diagram commutes

$$\begin{array}{ccccccc} \cdots & \longrightarrow & \Omega^{odd}(X) & \xrightarrow{d_\lambda} & \Omega^{even}(X) & \xrightarrow{d_\lambda} & \Omega^{odd}(X) \xrightarrow{d_\lambda} \cdots \\ & & \downarrow \psi_{2p-1} & & \downarrow \psi_{2p} & & \downarrow \psi_{2p+1} \\ \cdots & \longrightarrow & \Omega[[z]]_\lambda(X)^{2p-1} & \xrightarrow{d_\lambda} & \Omega[[z]]_\lambda(X)^{2p} & \xrightarrow{d_\lambda} & \Omega[[z]]_\lambda(X)^{2p+1} \xrightarrow{d_\lambda} \cdots \end{array}$$

In fact, for $e = 0, 1$ and $\omega = \sum_{i=0}^{\infty} \omega^{e+2i}$ we define

$$\psi_{2p+e}(\omega) := \sum_{i=0}^{\lfloor \frac{p}{2} \rfloor} \frac{z^{p-i} \omega^{e+2i}}{(p-i)!} .$$

If $p > \dim(X)$, then ψ_p is an isomorphism. Therefore for large p the isomorphisms ψ_p induce embeddings $H^*(X; \lambda) \hookrightarrow H^*(X; G)$. In this way $H^*(X; G)$ is a replacement of $H^*(X; \lambda)$ with good functorial properties.

1.3.24 The definition of real cohomology of X twisted by a gerbe as

$$H^*(X; G) := H^*(G; \mathbb{R})$$

has a couple of additional interesting features.

- (1) First of all note that \mathbb{R} is a commutative ring. Therefore $H^*(X; G)$ has naturally the structure of a graded commutative ring. In the old picture this structure seems to be partially reflected by the product

$$H^*(X; a\lambda) \otimes H^*(X; b\lambda) \rightarrow H^*(X; (a+b)\lambda) .$$

- (2) One can replace \mathbb{R} by any other abelian group. In particular, one can define integral twisted cohomology by $H^*(X; G; \mathbb{Z}) := H(G; \mathbb{Z})$. This definition of an integral twisted cohomology proposes a solution to the question raised in the remark made in [1, Sec. 6]. Using the maps ψ_p introduced above we can identify the image of $H^*(X; G; \mathbb{Z}) \rightarrow H^*(X; G)$ as a lattice in $H^*(X; \lambda)$. The result depends on the choice of p , and in view of the denominators in the formula for ψ_p the position of lattice is not very obvious.
- (3) In the proof of Theorem 1.1 we construct a de Rham model for the cohomology of $H^*(G; \mathbb{R})$. Let $\Omega_G^{<p}$ be the sheaf of de Rham complexes truncated at $p-1$ and form the sheaf of Deligne complexes $\mathcal{H}(p-1)_G := (i^\sharp \mathbb{Z}_G \rightarrow \Omega_G^{<p})$, where $i^\sharp \mathbb{Z}_G$ sits in degree -1 . We can then define the real Deligne cohomology $\hat{H}^p(G; \mathbb{Z})$ of G as the $(p-1)$ -st hypercohomology of the complex $\mathcal{H}_G(p-1)$ (see [5] for a definition of Deligne cohomology for manifolds in a similar fashion).

1.4 Sheaf theory for smooth stacks

1.4.1 This subsection is the introduction to the sheaf theoretic part of the paper. We consider a smooth stack X . In order to define the notion of a sheaf on X we associate to X a Grothendieck site \mathbf{X} . A site consists of a category \mathbf{X} and the choice of covering families $\text{cov}_{\mathbf{X}}(U)$ for the objects $U \in \mathbf{X}$. Presheaves on \mathbf{X} are just contravariant set-valued functors on \mathbf{X} . A sheaf on \mathbf{X} is a presheaf which satisfies a descend condition with respect to the covering families.

1.4.2 We define the category \mathbf{X} as a full subcategory of the category of manifolds U over X such that the structure map $U \rightarrow X$ is smooth. The covering families of $U \rightarrow X$ are families of submersions over X whose union maps surjectively to U . Observe that the category of smooth manifolds can be considered as a site with the above mentioned choice of covering families. By the Yoneda embedding it maps to the two-category of smooth stacks. In Subsection 2 we consider this abstract situation. We consider a site \mathcal{S} , a two-category \mathcal{C} and a functor $z : \mathcal{S} \rightarrow \mathcal{C}$. Furthermore we consider a subcategory $r\mathcal{C}$ which plays the role of the subcategory of stacks with smooth representable morphisms. In this situation we associate to each object $X \in \mathcal{C}$ the site \mathbf{X} (see 2.1) as the full subcategory of $(z(U) \rightarrow X) \in \mathcal{S}/X$ such that the structure map belongs to $r\mathcal{C}$. The covering families are induced from \mathcal{S} (see 2.2).

1.4.3 The central topic of Subsection 2 is the adjoint pair (2.5) of functors

$$f^* : \text{Pr } \mathbf{X} \Leftrightarrow \text{Pr } \mathbf{G} : f_*$$

between presheaf categories associated to a morphism $f : G \rightarrow X$. Since in general f does not induce a morphism between the sites \mathbf{G} and \mathbf{X} we define these functors in an ad-hoc manner (see 2.3 and 2.4). For two composable morphisms f, g we relate $(g \circ f)^*$, $(g \circ f)_*$ with $g_* \circ f_*$, $f^* \circ g^*$ in Lemma 2.6.

1.4.4 In Subsection 2.2 we specialize to smooth stacks. If the morphism $f : G \rightarrow H$ between smooth stacks is smooth or representable, then it gives rise to a morphism of sites f_{\sharp} or f^{\sharp} , respectively (2.2.6 and 2.2.7). We verify that our ad-hoc definitions of f^* or f_* , respectively, coincide with the standard functors induced from the morphism of sites f^{\sharp} or f_{\sharp} (see Lemmas 2.7, 2.9, 2.11).

1.4.5 Most of the statements which we formulate for the sheaf theory on stacks are well-known in the usual sheaf theory on sites and for functors associated to morphisms of sites. But for the sheaf theory on stacks we must be very careful about which of these standard facts remain true in general. For other statements we must know and under which additional assumptions they carry over to stacks.

1.4.6 An important point is the observation that for every morphism between smooth stacks the functor f_* preserves sheaves (Lemma 2.13). In the Lemmas 2.16 and 2.22 we study the compatibility of the pull back with the push forward in cartesian squares. In Lemma 2.23 we study under which additional assumptions we have relations like $(g \circ f)^* \cong f^* \circ g^*$.

1.4.7 In order to define the cohomology of a gerbe we must descend the functors f_* and f^* to the derived categories of presheaves and sheaves of abelian groups. This question is studied in Subsection 2.3. Here the exactness properties of the functors studied in the preceding subsections play an important role. Most of the statements in this subsection are standard for the usual sheaf theory and functors associated to a morphism of sites. Here we study carefully under which additional conditions they remain true for stacks.

1.4.8 The main result (Lemma 2.41) of Subsection 2.4 is that the derived functor Rf_* for a map $G \rightarrow X$ of smooth stack can be calculated using a simplicial approximation of $G \rightarrow X$. In particular, if X is a manifold, then the calculation of Rf_* can be reduced to ordinary sheaf theory on manifolds. We use this simplicial model in the proof of our main theorem, for the explicit calculation of the cohomology of the stack $[*/S^1]$ in Lemma 3.8, but also to verify that pull-back and push-forward commute on the level of derived functors for certain cartesian diagrams in Lemma 2.43.

1.4.9 The covering families of the small site (U) of a manifold are coverings by open subsets. Thus the sheaf theory for (U) is the ordinary one. If $(U \rightarrow X) \in \mathbf{X}$, then a presheaf on X induces a presheaf on (U) . In the present paper the sheaf theory on (U) is considered to be well-understood. The main goal of Subsection 2.5 is to compare the sheafification functors on \mathbf{X} and (U) (see Lemma 2.47). This result is very useful in explicit calculations since it says that certain questions can be studied for each $(U \rightarrow X) \in \mathbf{X}$ separately and with respect to the small site (U) . This sort of reasoning will be applied in the proof that the de Rham complex of a stack is a flabby resolution of the constant sheaf with value \mathbb{R} , where we use that this fact is well-known on each manifold equipped with the site (U) . It is also used in the proof of Lemma 2.50 which says that for a smooth map between smooth stacks the pull back commutes with the sheafification functor.

2 Sheaf theory for smooth stacks

2.1 Over sites

2.1.1 The goal of the present subsection is to develop some elements of sheaf theory in the following situation. Let \mathcal{S} be a site (see [24, 1.2.1] for a definition), \mathcal{C} a two-category with invertible two-morphisms, and $z : \mathcal{S} \rightarrow \mathcal{C}$ a functor (we consider \mathcal{S} as a two-category with only identity two-isomorphisms). Finally let $r\mathcal{C}$ be a subcategory which we call the category of admissible morphisms.

To each object $G \in \mathcal{C}$ we will associate a site \mathbf{G} (sometimes we will write $\text{Site}(G) := \mathbf{G}$) and the categories of presheaves $\text{Pr } \mathbf{G}$ and sheaves $\text{Sh } \mathbf{G}$ of sets on this site. For a morphism $f \in \mathcal{C}(G, H)$ we will define an adjoint pair of functors

$$f^* : \text{Pr } \mathbf{H} \Leftrightarrow \text{Pr } \mathbf{G} : f_* .$$

In general these functors are not induced by a morphism of sites.

2.1.2 Let $G \in \mathcal{C}$. We define the underlying category of \mathbf{G} .

Definition 2.1 *The objects of \mathbf{G} are pairs (U, ϕ) , where $U \in \mathcal{S}$ and $\phi \in r\mathcal{C}(z(U), G)$. A morphism $(U, \phi) \rightarrow (U', \phi')$ is given by a pair (h, σ) , where $h \in \mathcal{S}(U, U')$ and σ is a two-isomorphism*

$$\begin{array}{ccc} z(U) & \xrightarrow{z(h)} & z(U') \\ & \searrow \phi & \downarrow \sigma & \swarrow \phi' \\ & & G & \end{array}$$

The composition in \mathbf{G} is defined in the obvious way.

Sometimes we will abbreviate the notation and write U or $(z(U) \rightarrow G)$ for (U, ϕ) .

2.1.3 Next we define the coverings of an object (U, ϕ) of \mathbf{G} .

Definition 2.2 *A covering of (U, ϕ) is a collection of morphisms*

$$((U_i, \phi_i) \xrightarrow{(h_i, \sigma_i)} (U, \phi))_{i \in I}$$

such that $(U_i \xrightarrow{h_i} U)_{i \in I}$ is a covering of U in \mathcal{S} .

In fact it is easy to verify the axioms listed in the definition [24, 1.2.1]. The only non-obvious part asserts that given a covering $((U_i, \phi_i) \rightarrow (U, \phi))_{i \in I}$ and a morphism $(V, \psi) \rightarrow (U, \phi)$, then the fibre products $(V_i, \psi_i) := (V, \psi) \times_{(U, \phi)} (U_i, \phi_i)$ exist in \mathbf{G} and $((V_i, \psi_i) \rightarrow (V, \psi))_{i \in I}$ is a covering of (V, ψ) . By a little diagram chase one verifies that $(V, \psi) \times_{(U, \phi)} (U_i, \phi_i) \cong (V \times_U U_i, \phi \circ z(\kappa))$, where $V \times_U U_i$ is the fibre product in \mathcal{S} and $\kappa : V \times_U U_i \rightarrow U$ is the natural map.

2.1.4 Let $f : G \rightarrow H$ be a morphism in \mathcal{C} . Then we can define the functor $f^* : \text{Pr } \mathbf{H} \rightarrow \text{Pr } \mathbf{G}$ as follows. Given $(z(V) \rightarrow G) \in \mathbf{G}$ we consider the category V/\mathbf{H} (recall that V abbreviates $(z(V) \rightarrow G)$) of diagrams

$$\begin{array}{ccc} V & & z(V) \longrightarrow G \\ \downarrow & & \downarrow \quad \nearrow \quad \downarrow f \\ U & & z(U) \longrightarrow H \end{array} .$$

A morphism in this category is given by a morphism $(z(U') \rightarrow H) \rightarrow (z(U) \rightarrow H)$ in \mathbf{H} fitting into

$$\begin{array}{ccc}
 & & U \\
 & \nearrow & \uparrow \\
 V & & \\
 & \searrow & \downarrow \\
 & & U'
 \end{array}
 \quad
 \begin{array}{ccc}
 z(V) & \longrightarrow & G \\
 \searrow & \nearrow & \downarrow f \\
 & z(U) & \longrightarrow H \\
 \nearrow & \downarrow & \parallel \\
 z(U') & \longrightarrow & H
 \end{array}
 .$$

Let $F \in \text{Pr } \mathbf{H}$.

Definition 2.3 *We define*

$$f^*F(V) := \text{colim}_{V/\mathbf{H}} F(U) .$$

A morphism $V' \rightarrow V$ in \mathbf{G} induces naturally a functor $V/\mathbf{H} \rightarrow V'/\mathbf{H}$. The relevant diagram is

$$\begin{array}{ccc}
 V' & & z(V') \longrightarrow G \\
 \downarrow & & \searrow \nearrow \\
 V & & z(V) \longrightarrow G \\
 \downarrow & & \searrow \nearrow \\
 U & & z(U) \longrightarrow H
 \end{array}
 \quad
 \begin{array}{ccc}
 & & G \\
 & \nearrow & \parallel \\
 & z(V) & \longrightarrow G \\
 & \searrow & \downarrow f \\
 & z(U) & \longrightarrow H
 \end{array}
 .$$

We therefore get a map $f^*F(V) \rightarrow f^*F(V')$, and this makes f^*F a presheaf on \mathbf{G} .

2.1.5 Let $f : G \rightarrow H$ again be a morphism in \mathcal{C} . We define a functor $f_* : \text{Pr } \mathbf{G} \rightarrow \text{Pr } \mathbf{H}$ as follows. We consider $(z(U) \rightarrow G) \in \mathbf{H}$. Then we consider the category of diagrams \mathbf{G}/U of diagrams

$$\begin{array}{ccc}
 V & & z(V) \longrightarrow G \\
 \downarrow & & \searrow \nearrow \\
 U & & z(U) \longrightarrow H
 \end{array}
 \quad
 \begin{array}{ccc}
 & & G \\
 & \nearrow & \downarrow f \\
 & z(V) & \longrightarrow G \\
 & \searrow & \downarrow \\
 & z(U) & \longrightarrow H
 \end{array}
 .$$

A morphism of such diagrams is given by a morphism $V' \rightarrow V$ in \mathbf{G} which fits into

$$\begin{array}{ccc}
 V' & & z(V') \longrightarrow G \\
 \downarrow & & \searrow \nearrow \\
 V & & z(V) \longrightarrow G \\
 \downarrow & & \searrow \nearrow \\
 U & & z(U) \longrightarrow H
 \end{array}
 \quad
 \begin{array}{ccc}
 & & G \\
 & \nearrow & \parallel \\
 & z(V) & \longrightarrow G \\
 & \searrow & \downarrow f \\
 & z(U) & \longrightarrow H
 \end{array}
 .$$

Definition 2.4 *We define*

$$f_*F(U) := \lim_{\mathbf{G}/U} F(V) .$$

A morphism $U \rightarrow U'$ in \mathbf{H} induces naturally a functor $\mathbf{G}/U \rightarrow \mathbf{G}/U'$. The relevant diagram is

$$\begin{array}{ccc}
 V & & z(V) \longrightarrow G \\
 \downarrow & & \downarrow \nearrow \\
 U & & z(U) \longrightarrow G \\
 \downarrow & & \downarrow \nearrow \\
 U' & & z(U') \longrightarrow H
 \end{array}
 \begin{array}{c}
 \\
 \\
 \\
 \\
 \downarrow f \\
 \\
 \\
 \\
 \downarrow f
 \end{array}
 .$$

We therefore get a map $f_*F(U') \rightarrow f_*F(U)$, and this makes f_*F a presheaf on \mathbf{H} .

2.1.6 Let $f \in \mathcal{C}(G, H)$ as before.

Lemma 2.5 *The functors f_* and f^* naturally form an adjoint pair*

$$f^* : \text{Pr } \mathbf{H} \Leftrightarrow \text{Pr } \mathbf{G} : f_* .$$

Proof. We give the unit and the counit. Let $(z(W) \rightarrow G) \in \mathbf{G}$. Then

$$f^* f_* F(W) = \text{colim}_U \lim_V F(V) ,$$

where the colimit-limit is taken over a category of diagrams

$$\begin{array}{ccc}
 V & & z(V) \longrightarrow G \\
 \downarrow & & \downarrow \\
 U & & z(U) \longrightarrow H \\
 \uparrow & & \uparrow \\
 W & & z(W) \longrightarrow G
 \end{array}
 \begin{array}{c}
 \\
 \\
 \\
 \\
 \downarrow f \\
 \\
 \uparrow f \\
 \\
 \downarrow f
 \end{array}$$

(we leave out the two-isomorphisms). The counit is a natural transformation

$$f^* f_* F(W) \rightarrow F(W) .$$

It is given by the universal property of the colimit and the collection of maps which associates to U the canonical map $\lim_V F(V) \rightarrow F(W)$.

Furthermore, let $(z(U) \rightarrow H) \in \mathbf{H}$. Then

$$f_* f^* F(U) = \lim_V \text{colim}_W F(W) ,$$

where the limit-colimit is taken over a category of diagrams

$$\begin{array}{ccc}
 U & & z(U) \longrightarrow H \\
 \uparrow & & \uparrow \\
 V & & z(V) \longrightarrow G \\
 \downarrow & & \downarrow \\
 W & & z(W) \longrightarrow H
 \end{array}
 \begin{array}{c}
 \\
 \uparrow f \\
 \\
 \downarrow f \\
 \\
 \uparrow f \\
 \\
 \downarrow f
 \end{array}$$

(we leave out the two-isomorphisms). The unit is a natural transformation

$$F(U) \rightarrow f_* f^* F(U) .$$

It is given by the universal property of the limit and the collection of maps which associates to V the natural map $F(U) \rightarrow \operatorname{colim}_W F(W)$.

We leave it to the interested reader to perform the remaining checks. \square

2.1.7 Let us consider a pair of composable maps in \mathcal{C}

$$G \xrightarrow{f} H \xrightarrow{g} L .$$

Lemma 2.6 *We have natural transformations of functors*

$$(g \circ f)_* \rightarrow g_* \circ f_* , \quad f^* \circ g^* \rightarrow (g \circ f)^* .$$

Proof. We discuss the transformation $f^* \circ g^* \rightarrow (g \circ f)^*$. Let $F \in \operatorname{Pr} \mathbf{L}$ and $(z(W) \rightarrow G) \in \mathbf{G}$. Inserting the definitions we have

$$f^* \circ g^*(F)(W) \cong \operatorname{colim}_{\mathcal{A}} F(U) ,$$

where \mathcal{A} is the category of diagrams

$$\begin{array}{ccc} z(W) & \longrightarrow & G \\ \downarrow & \nearrow \text{dotted} & \downarrow f \\ z(V) & \longrightarrow & H \\ \downarrow & \nearrow \text{dotted} & \downarrow g \\ z(U) & \longrightarrow & L \end{array}$$

with $(V \rightarrow H) \in \mathbf{H}$ and $(U \rightarrow L) \in \mathbf{L}$. The vertical composition provides a functor $\mathcal{A} \rightarrow W/\mathbf{L}$, where W/\mathbf{L} is the category of diagrams of the form

$$\begin{array}{ccc} z(W) & \longrightarrow & G \\ \downarrow & \nearrow \text{dotted} & \downarrow g \circ f \\ z(U) & \longrightarrow & L \end{array} .$$

We get an induced map of colimits

$$f^* \circ g^*(F)(W) \rightarrow (g \circ f)^* F(W) = \operatorname{colim}_{W/\mathbf{L}} F(U) .$$

The other transformation $(g \circ f)_* \rightarrow g_* \circ f_*$ is obtained in a similar manner or, equivalently, by adjointness. \square

In general, we can not expect that these transformations are isomorphisms. But under additional assumptions they are, see 2.23.

2.2 The site of a smooth stack

2.2.1 We consider the small² site \mathbf{Mf}^∞ of smooth manifolds. Its underlying category is the category of smooth manifolds and smooth maps. A collection of smooth maps $(U_i \rightarrow U)_{i \in I}$ is a covering if and only if this family is isomorphic to the collection of inclusions of the open subsets of U given by an open covering of U .

We use the small site \mathbf{Mf}^∞ in order to define stacks in smooth manifolds. We refer to [10], [19], [22] for the language of stacks.

²In the present paper the adjectives *small* and *big* refer to the type of covering families. In fact, in order to avoid set-theoretic problems one must require that a site is a small category. In the present paper we will ignore this problem. It can be resolved by either working with universes or replacing \mathbf{Mf}^∞ by an equivalent small category (see e.g. [19]).

2.2.2 We will also consider the big site \mathcal{S} on smooth manifolds. In this site a family $(U_i \rightarrow U)_{i \in I}$ of smooth maps is a covering if the maps $U_i \rightarrow U$ are submersions and $\sqcup_{i \in I} U_i \rightarrow U$ is surjective. We will use this site in order to define the site of a stack according to 2.1.

2.2.3 In this paragraph we recall the main notions of the theory of smooth stacks.

- (1) A morphism of stacks $G \rightarrow H$ is called representable, if for each manifold U and map $U \rightarrow H$ the fibre product $U \times_H G$ is equivalent to a manifold. A composition of representable maps is representable.
- (2) A representable morphism $G \rightarrow H$ of stacks is called smooth if for each manifold U and map $U \rightarrow H$ the induced map $U \times_H G \rightarrow U$ (of manifolds) is a submersion.
- (3) A map $U \rightarrow G$ from a manifold to a stack is called an atlas if it is representable, smooth and admits local sections.
- (4) A stack in smooth manifolds is called smooth if it admits an atlas [10, Def. 2.4].
- (5) A morphism (not necessarily representable) between smooth stacks $f : G \rightarrow H$ is called smooth if for an atlas $A \rightarrow G$ the composition $A \rightarrow G \rightarrow H$ is smooth [10, Def. 2.10]. A composition of smooth maps is smooth.
- (6) A smooth morphism $U \rightarrow G$ from a manifold to a smooth stack is representable.

2.2.4 Let \mathcal{C} be the two-category of smooth stacks in smooth manifolds. We have a Yoneda embedding $z : \mathcal{S} \rightarrow \mathcal{C}$. Note that in general we will omit the Yoneda embedding in the notation and consider \mathcal{S} as a subcategory of \mathcal{C} . We let $r\mathcal{C}$ be the subcategory of representable smooth morphisms.

2.2.5 The conventions introduced in 2.2.4 place us in the situation of 2.1.1. Let $G \in \mathcal{C}$ be a smooth stack. Then by \mathbf{G} we denote the site according to Definitions 2.1 and 2.2. Note that this site is derived from the big site \mathcal{S} on smooth manifolds. We now have the categories of presheaves $\text{Pr } \mathbf{G}$ and sheaves $\text{Sh } \mathbf{G}$ on the stack G . The present definition is essentially the same as in [10, Sec. 4] except that we use the big site \mathcal{S} on the category of smooth manifolds instead of the small site Mf^∞ . Note that we use the descend property for the big site in the proof of Lemma 2.36. The relation between the sheaf theories on the big and the small sites will be discussed in [6].

2.2.6 Let $f : G \rightarrow H$ be a representable morphism of smooth stacks in smooth manifolds. Then it induces a morphism of sites $f^\sharp : \mathbf{H} \rightarrow \mathbf{G}$ by the rule $f^\sharp(U \rightarrow H) := U \times_H G \rightarrow G$ (it is easy to check the axioms listed in [24, 1.2.2]).

2.2.7 If $f : G \rightarrow H$ is a smooth morphism of smooth stacks in smooth manifolds, then we can define another morphism of sites $f_\sharp : \mathbf{G} \rightarrow \mathbf{H}$ by $f_\sharp(V \rightarrow G) := (V \rightarrow G \xrightarrow{f} H)$.

2.2.8 We call a functor left exact if it preserves arbitrary limits. If it preserves arbitrary colimits, then we call it right exact. A functor is said to be exact if it is right and left exact.

Recall that a functor which is a left adjoint is right exact. Similarly, a right adjoint is left exact.

2.2.9 A morphism of sites $q : \mathbf{H} \rightarrow \mathbf{G}$ induces an adjoint pair

$$q_* : \text{Pr } \mathbf{H} \Leftrightarrow \text{Pr } \mathbf{G} : q^* .$$

(see [24, 2.3]). In the following we compare these maps with the ad-hoc definitions 2.3 and 2.4 and discuss some special properties.

2.2.10

Lemma 2.7 *If $f : G \rightarrow H$ is a smooth morphism between smooth stacks, then we have $f^* \cong (f_\sharp)^*$. In particular, then f^* is exact and preserves sheaves.*

Proof. Let $(V \rightarrow G) \in \mathbf{G}$. According to the definition [24, 2.3] we have

$$(f_\sharp)^*(F)(V \rightarrow G) := F(f_\sharp(V \rightarrow G)) = F(V \rightarrow G \rightarrow H) .$$

If $(V \rightarrow G) \in \mathbf{G}$, then the category V/\mathbf{H} has an initial object

$$\begin{array}{ccc} V & \longrightarrow & G \\ \parallel & \nearrow \text{id} & \downarrow f \\ V & \longrightarrow & H \end{array} .$$

Therefore

$$(f^*F)(V \rightarrow G) \cong F(V \rightarrow G \rightarrow H) . \quad (2.8)$$

This implies that $f^* \cong (f^\sharp)^*$.

It is well-known [24, 3.6] that the contravariant functor (in our case $(f^\sharp)^*$) associated to a morphism of sites preserves sheaves. Therefore f^* preserves sheaves.

The limit of a diagram of presheaves is defined objectwise. By (2.8) the functor f^* commutes with limits. As a left adjoint (by Lemma 2.5) it also commutes with colimits. \square

2.2.11 Let $f : G \rightarrow H$ be a representable and smooth morphism of smooth stacks.

Lemma 2.9 *We have an isomorphism of functors $(f^\sharp)_* \cong f^*$.*

Proof. Let $F \in \text{Pr } \mathbf{H}$. For $(V \rightarrow G) \in \mathbf{G}$ we have the category V/f^\sharp of pairs $((U \rightarrow H) \in \mathbf{H}, (V \rightarrow f^\sharp(U)) \in \text{Mor}(\mathbf{G}))$. It has a natural evaluation $\text{ev}_V : V/f^\sharp \rightarrow \mathbf{H}$ which maps $((U \rightarrow H), (V \rightarrow f^\sharp(U)))$ to $(U \rightarrow H)$. By definition (see [24, Proof of 2.3.1])

$$(f^\sharp)_*(F)(V) = \text{colim}_{V/f^\sharp} F \circ \text{ev}_V .$$

Now we observe that V/f^\sharp can be identified with the category of diagrams

$$\begin{array}{ccc} V & \longrightarrow & U \\ \downarrow & \nearrow & \downarrow \\ G & \xrightarrow{f} & H \end{array} .$$

Since f is smooth we see that $(V \rightarrow G \xrightarrow{f} H) \in \mathbf{H}$ and

$$\begin{array}{ccc} V & \xrightarrow{\text{id}_V} & V \\ \downarrow & \nearrow \text{id} & \downarrow \\ G & \xrightarrow{f} & H \end{array}$$

is the initial element of V/f^\sharp . We conclude that

$$(f^\sharp)_*(F)(V \rightarrow G) \cong F(V \rightarrow G \rightarrow H) . \quad (2.10)$$

The equality $f^* \cong (f^\sharp)_*$ now follows from (2.8). \square

One can not expect that f^* is left exact for a general map $f : G \rightarrow H$. In fact this problem occurs in the corresponding definition in [14] of the pull-back for the lisse-etale site of an algebraic stack. For more details and a solution see [23].

2.2.12

Lemma 2.11 *If $f : G \rightarrow H$ is a representable morphism of smooth stacks, then $f_* = (f^\sharp)^* : \text{Pr } \mathbf{G} \rightarrow \text{Pr } \mathbf{H}$. The functor f_* is exact.*

Proof. Let $(U \rightarrow H) \in \mathbf{H}$. Then $f^\sharp(U \rightarrow H) = (U \times_H G \rightarrow G)$ is the final object in \mathbf{G}/U . Therefore

$$f_*F(U) \cong F(U \times_H G) \cong (f^\sharp)^*F(U) . \quad (2.12)$$

Since $(f^\sharp)^*$ is a right adjoint it commutes with limits. Since colimits of presheaves are defined objectwise it follows from the formula (2.12) that f_* also commutes with colimits. \square

2.2.13 Let now $f : G \rightarrow H$ be a map of smooth stacks.

Lemma 2.13 *The functor f_* preserves sheaves.*

Proof. Let $F \in \mathbf{Sh}\mathbf{G}$. Consider $(U \rightarrow H) \in \mathbf{H}$ and let $(U_i \rightarrow U)$ be a covering of U . Consider a diagram

$$\begin{array}{ccc} V & \longrightarrow & G \\ \downarrow & \nearrow & \downarrow f \\ U & \longrightarrow & H \end{array} . \quad (2.14)$$

From this we obtain a collection of diagrams

$$\begin{array}{ccc} V_i := U_i \times_U V & \longrightarrow & G \\ \downarrow & \nearrow & \downarrow f \\ U_i & \longrightarrow & H \end{array}$$

functorially in V . Observe that $(V_i \rightarrow V)$ is a covering in \mathbf{G} . We now consider the map of diagrams

$$\begin{array}{ccccc} f_*F(U) & \longrightarrow & \prod_i f_*F(U_i) & \Longrightarrow & \prod_{i,j} f_*F(U_i \times_U U_j) \\ \downarrow & & \downarrow & & \downarrow \\ F(V) & \longrightarrow & \prod_i F(V_i) & \Longrightarrow & \prod_{i,j} F(V_i \times_V V_j) \end{array}$$

The vertical maps are given by specialization. We must show that the upper horizontal line is an equalizer diagram. The lower horizontal line has this property since F is a sheaf.

We now take the limit over the category of diagrams (2.14) and use the fact that a limit preserves equalizer diagrams. We get the commutative diagram of sets

$$\begin{array}{ccccc} f_*F(U) & \longrightarrow & \prod_i f_*F(U_i) & \Longrightarrow & \prod_{i,j} f_*F(U_i \times_U U_j) \\ \parallel & & \downarrow s & & \downarrow \\ f_*F(U) & \longrightarrow & \lim \prod_i F(V_i) & \Longrightarrow & \lim \prod_{i,j} F(V_i \times_V V_j) \end{array}$$

Let us assume that s is injective. Then the fact that the lower horizontal line is an equalizer diagram implies by a simple diagram chase that the upper horizontal line is an equalizer diagram.

We now show that s is injective. Note that a priori the product of specialization maps

$$s = \prod_i s_i : \prod_i f_*F(U_i) \rightarrow \prod_i \lim F(V_i)$$

may not be injective since the functors $L_i : \mathbf{G}/U \ni V \mapsto V_i \in \mathbf{G}/U_i$ are not necessarily essentially surjective. But in our situation the maps s_i are injective since each object in \mathbf{G}/U_i maps into an object in the image of L_i . To see this consider a diagram

$$\begin{array}{ccc} W & \xrightarrow{t} & G \\ \downarrow & \nearrow & \downarrow f \\ U_i & \longrightarrow & H \end{array} \in \mathbf{G}/U_i .$$

Using the composition $W \rightarrow U_i \rightarrow U$ we can form the diagram

$$\begin{array}{ccc} W & \longrightarrow & G \in \mathbf{G}/U \\ \downarrow & \nearrow & \downarrow f \\ U & \longrightarrow & X \end{array}$$

and define the morphism in \mathbf{G}/U_i

$$\begin{array}{ccc} & U_i \times_U W & \\ \text{pr}_1 \curvearrowright & \uparrow j & \searrow \text{topr}_2 \\ & W & \longrightarrow G \\ & \downarrow & \nearrow \\ & U_i & \longrightarrow H \end{array}$$

where $j : W \rightarrow U_i \times_U W$ is induced by $W \rightarrow U_i$ and $\text{id}_W : W \rightarrow W$.

□

2.2.14 Assume that we have a diagram in smooth stacks

$$\begin{array}{ccc} G & \xrightarrow{u} & H \\ \downarrow f & \nearrow & \downarrow g \\ M & \xrightarrow{v} & N \end{array}, \tag{2.15}$$

where u and v are smooth.

Lemma 2.16 *We have a natural map of functors $\text{Pr } \mathbf{H} \rightarrow \text{Pr } \mathbf{M}$*

$$v^* \circ g_* \rightarrow f_* \circ u^*$$

which is an isomorphism if (2.15) is cartesian.

Proof. We use the description (2.8) of v^* obtained in the proof of Lemma 2.7. Let $(U \rightarrow M) \in \mathbf{M}$. Then we have

$$v^* \circ g_* F(U) \cong \lim F(A),$$

where the limit is taken over a category D of diagrams

$$\begin{array}{ccccc} & & & & A \\ & & & \swarrow & \downarrow \\ & G & \xrightarrow{u} & H & \\ & \downarrow f & \nearrow & \downarrow g & \\ & M & \xrightarrow{v} & N & \\ & \swarrow & \nearrow & \swarrow \sigma & \downarrow \\ U & & & & U \end{array}$$

(2.17)

(where A varies). On the other hand $f_* \circ u^*(F)(U) \cong \lim F(V)$, where the limit is taken over the category E of diagrams

$$\begin{array}{ccc}
 V & \xrightarrow{\quad} & V \\
 \downarrow & \searrow & \swarrow \\
 & G & \xrightarrow{u} & H \\
 & \downarrow f & \searrow & \downarrow g \\
 & M & \xrightarrow{v} & N \\
 \uparrow & \swarrow & \nwarrow & \uparrow \\
 U & & & U
 \end{array}
 \tag{2.18}$$

(V varies). We define a functor $X : E \rightarrow D$ which sends the diagram (2.18) to the diagram

$$\begin{array}{ccc}
 & & V \\
 & & \downarrow \\
 & G & \xrightarrow{u} & H \\
 & \downarrow f & \searrow & \downarrow g \\
 & M & \xrightarrow{v} & N \\
 \uparrow & \swarrow & \nwarrow & \uparrow \\
 U & \xrightarrow{\text{id}} & U
 \end{array}$$

We write F_E and F_D for the functor F precomposed with the evaluations $E \rightarrow \mathbf{H}$ and $D \rightarrow \mathbf{H}$. The identity $F(V) \xrightarrow{\sim} F(V)$ induces an isomorphism $F_D \circ X \xrightarrow{\sim} F_E$. Therefore we have a natural map of limits

$$v^* \circ g_* F(U) \rightarrow f_* \circ u^* F(U) .
 \tag{2.19}$$

This gives the required transformation of functors.

If (2.15) is cartesian, then we can define a functor $Y : D \rightarrow E$ which maps the diagram (2.17) functorially to

$$\begin{array}{ccc}
 U \times_N A & \xrightarrow{h} & G & \xrightarrow{u} & H \\
 \downarrow & \searrow & \downarrow f & \searrow & \downarrow g \\
 & M & \xrightarrow{v} & N \\
 \uparrow & \swarrow & \nwarrow & \uparrow \\
 U & \xrightarrow{\text{id}} & U
 \end{array}$$

which employs the map $A \rightarrow H \rightarrow N$. The map h is induced by the universal property of the cartesian diagram. Since $U \rightarrow M$ and $A \rightarrow H$ are smooth, the map h is smooth, too. The map $A \rightarrow U$ together with the two-isomorphism σ gives a map $A \rightarrow U \times_N A$ in \mathbf{H} . This map induces the natural transformation $F_E \circ Y \rightarrow F_D$. It gives a map of limits

$$f_* \circ u^*(F)(U) \rightarrow v^* \circ g_* F(U) .
 \tag{2.20}$$

One can check that (2.20) is inverse to (2.19). □

2.2.15 Assume again that we have a diagram in smooth stacks

$$\begin{array}{ccc} G & \xrightarrow{u} & H \\ \downarrow f & \nearrow & \downarrow g \\ M & \xrightarrow{v} & N \end{array} . \quad (2.21)$$

We now assume that f and g are representable, and that u, v are smooth.

Lemma 2.22 *We have a natural map of functors $\mathrm{Pr} \mathbf{H} \rightarrow \mathrm{Pr} \mathbf{M}$*

$$v^* \circ g_* \rightarrow f_* \circ u^*$$

which is an isomorphism if (2.21) is cartesian.

This is a special case of Lemma 2.16. But under the additional representability assumptions on f and g the proof simplifies considerably.

Proof. Let $F \in \mathrm{Pr} \mathbf{H}$. For $(U \rightarrow M) \in \mathbf{M}$ we calculate

$$\begin{aligned} v^* \circ g_*(F)(U) &\cong \operatorname{colim}_{V \in U/\mathbf{N}} \lim_{W \in \mathbf{H}/V} F(W) \\ &\stackrel{(2.8)}{\cong} \lim_{W \in \mathbf{H}/(U \rightarrow M \rightarrow N)} F(W) \\ &\stackrel{(2.12)}{\cong} F(H \times_N U \rightarrow H) . \end{aligned}$$

On the other hand

$$\begin{aligned} f_* \circ u^*(F)(U) &\cong \lim_{Z \in \mathbf{G}/U} \operatorname{colim}_{W \in Z/\mathbf{H}} F(W) \\ &\stackrel{(2.12)}{\cong} \operatorname{colim}_{W \in (U \times_M G)/\mathbf{H}} F(W) \\ &\stackrel{(2.8)}{\cong} F(U \times_M G \rightarrow G \rightarrow H) . \end{aligned}$$

The transformation $v^* \circ g_*(F)(U) \rightarrow f_* \circ u^*(F)(U)$ is now induced from the map $(G \times_M U \rightarrow H \times_N U) \in \mathbf{H}$. If the diagram is cartesian, then we have $G \times_M U \cong (H \times_N M) \times_M U \cong H \times_N U$ so that the transformation is an isomorphism. \square

2.2.16 Let us consider a pair of composable maps of smooth stacks

$$G \xrightarrow{f} H \xrightarrow{g} L .$$

In Lemma 2.6 we have found natural transformations of functors between presheaf categories

$$(g \circ f)_* \rightarrow g_* \circ f_* , \quad f^* \circ g^* \rightarrow (g \circ f)^* .$$

Lemma 2.23 *If g is representable, or if f is smooth, then these transformations are isomorphisms.*

Proof. We consider the transformation $f^* \circ g^* \rightarrow (g \circ f)^*$ which appears as a transformation of colimits induced by a functor between indexing categories $\mathcal{A} \rightarrow W/\mathbf{L}$, where we use the notation introduced in the proof of Lemma 2.6. Under the present additional assumptions on f or g we have a functor $W/\mathbf{L} \rightarrow \mathcal{A}$ which induces the inverse of the transformation. In the following we describe these functors.

If g is representable, then each diagram

$$\begin{array}{ccc} W & \longrightarrow & G \\ \downarrow & \nearrow & \downarrow g \circ f \\ U & \longrightarrow & L \end{array} \quad (2.24)$$

in W/\mathbf{L} naturally completes to

$$\begin{array}{ccc} W & \longrightarrow & G \\ \downarrow & \nearrow & \downarrow f \\ U \times_L H & \longrightarrow & H \\ \downarrow & \nearrow & \downarrow g \\ U & \longrightarrow & L \end{array}$$

in \mathcal{A} .

If f is smooth, then the diagram (2.24) can be naturally completed to

$$\begin{array}{ccc} W & \longrightarrow & G \\ \parallel & \nearrow \text{id} & \downarrow f \\ W & \longrightarrow & H \\ \downarrow & \nearrow & \downarrow g \\ U & \longrightarrow & L \end{array}$$

in \mathcal{A} .

It follows from adjointness that under the additional assumptions on f or g the transformation $(g \circ f)_* \rightarrow g_* \circ f_*$ is an isomorphism, too. \square

2.2.17 Let $f : G \rightarrow H$ be a smooth map of smooth stacks.

Lemma 2.25 *There exists a functor $f_! : \text{Pr } \mathbf{G} \rightarrow \text{Pr } \mathbf{H}$ so that we get an adjoint pair*

$$f_! : \text{Pr } \mathbf{G} \Leftrightarrow \text{Pr } \mathbf{H} : f^* .$$

Proof. Let $(V \rightarrow G) \in \mathbf{G}$. Then by (2.8) we have $f^*F(V) \cong F(V \rightarrow G \rightarrow H)$.

Let $(V \rightarrow G) \in \mathbf{G}$ and $h_{V \rightarrow G} \in \text{Pr } \mathbf{G}$ be the corresponding representable presheaf. Then we have a natural isomorphism

$$\begin{aligned} \text{Hom}(h_{V \rightarrow G}, f^*F) &\cong f^*F(V \rightarrow G) \\ &\cong F(V \rightarrow G \rightarrow H) \\ &\cong \text{Hom}(h_{V \rightarrow G \rightarrow H}, F) \end{aligned}$$

which leads us to the definition

$$f_!h_{V \rightarrow G} := h_{V \rightarrow G \rightarrow H} .$$

If $L \in \text{Pr } \mathbf{G}$, then we can write $L \cong \text{colim}_{h_{V \rightarrow G} \rightarrow L} h_{V \rightarrow G}$. Since a left-adjoint must commute with colimits we are forced to set

$$f_!L := \text{colim}_{h_{V \rightarrow G} \rightarrow L} h_{V \rightarrow G \rightarrow H} .$$

Then we have indeed

$$\begin{aligned}
\mathrm{Hom}(L, f^*F) &\cong \mathrm{Hom}(\mathrm{colim}_{h_{V \rightarrow G} \rightarrow L} h_{V \rightarrow G}, f^*F) \\
&\cong \lim_{h_{V \rightarrow G} \rightarrow L} \mathrm{Hom}(h_{V \rightarrow G}, f^*F) \\
&\cong \lim_{h_{V \rightarrow G} \rightarrow L} \mathrm{Hom}(h_{V \rightarrow G \rightarrow H}, F) \\
&\cong \mathrm{Hom}(\mathrm{colim}_{h_{V \rightarrow G} \rightarrow L} h_{V \rightarrow G \rightarrow H}, F) \\
&\cong \mathrm{Hom}(f_!L, F)
\end{aligned}$$

□

2.3 Presheaves of abelian groups and derived functors

2.3.1 In the previous Subsections 2.1 and 2.2 we have developed a theory of set-valued presheaves and sheaves on stacks. We are in particular interested in the abelian categories of presheaves and sheaves of abelian groups and their derived categories. The functors (f^*, f_*) and (i^\sharp, i) preserve abelian group valued objects. In the present subsection we study how these functors descend to the derived categories. Furthermore, we check some functorial properties of these descended functors which will be employed in later calculations.

The derived version (Lemma 2.43) of the fact that pull-back commutes with push-forward in certain cartesian diagrams (Lemma 2.16) would fit into the present subsection, but can only be shown after the development of a computational tool in Subsection 2.4.

A similar remark applies to Lemma 2.50 saying that sheafification commutes with pull-back along smooth maps between smooth stacks. We will show this Lemma in Subsection 2.5.

2.3.2 For a site \mathbf{G} let $\mathrm{Pr}_{\mathrm{Ab}} \mathbf{G}$ and $\mathrm{Sh}_{\mathrm{Ab}} \mathbf{G}$ denote the abelian categories of presheaves and sheaves of abelian groups on \mathbf{G} . These categories have enough injectives [24, 2.1.1 and 2.1.2]. Let $D^+(\mathrm{Pr}_{\mathrm{Ab}} \mathbf{G})$ and $D^+(\mathrm{Sh}_{\mathrm{Ab}} \mathbf{G})$ denote the lower bounded derived categories of $\mathrm{Pr}_{\mathrm{Ab}} \mathbf{G}$ and $\mathrm{Sh}_{\mathrm{Ab}} \mathbf{G}$.

2.3.3 If $f : G \rightarrow H$ is a morphism of smooth stacks then $f_* : \mathrm{Pr} \mathbf{G} \rightarrow \mathrm{Pr} \mathbf{H}$ is left exact since it is a right adjoint. We therefore have the right derived functor $Rf_* : D^+(\mathrm{Pr}_{\mathrm{Ab}} \mathbf{G}) \rightarrow D^+(\mathrm{Pr}_{\mathrm{Ab}} \mathbf{H})$.

If $g : H \rightarrow L$ is a second morphism of smooth stacks, then we have a natural transformation

$$R(g \circ f)_* \rightarrow Rg_* \circ Rf_* .$$

In fact, let $F \in D^+(\mathrm{Pr}_{\mathrm{Ab}} \mathbf{G})$ be a lower bounded complex of injective presheaves. Then we choose an injective resolution $f_*F \rightarrow J$. Note that $g_*(J)$ represents $Rg_* \circ Rf_*(F)$. Then using (2.6) the required morphism is defined as the composition

$$R(g \circ f)_*(F) \cong (g \circ f)_*(F) \rightarrow g_* \circ f_*(F) \rightarrow g_*(J) \cong Rg_* \circ Rf_*(F) .$$

Lemma 2.26 *If f is smooth or g is representable, then*

$$R(g \circ f)_* \cong Rg_* \circ Rf_* .$$

Proof. If f is smooth, then f^* is exact. In this case f_* preserves injectives and we can take $J := f_*(F)$. We can now apply Lemma 2.23 in order to see that the natural transformation $(g \circ f)_*(F) \rightarrow g_* \circ f_*(F)$ is an isomorphism.

If g is representable, then g_* is exact by Lemma 2.11. In this case we have again by Lemma 2.23 that $R(g \circ f)_*(F) \cong (g \circ f)_*(F) \cong g_* \circ f_*(F) \cong Rg_* \circ Rf_*(F)$. □

2.3.4 Let $i : \mathbf{Sh}\mathbf{G} \rightarrow \mathbf{Pr}\mathbf{G}$ denote the inclusion. It has a left adjoint $i^\sharp : \mathbf{Pr}\mathbf{G} \rightarrow \mathbf{Sh}\mathbf{G}$, the sheafification functor (see [24, 3.1.1, 3.2.1]). Since the functor i is a right adjoint, it is left exact. We can form its right derived $Ri : D^+(\mathbf{Sh}_{\mathbf{Ab}}\mathbf{G}) \rightarrow D^+(\mathbf{Pr}_{\mathbf{Ab}}\mathbf{G})$.

Let $f : G \rightarrow H$ be a morphism of smooth stacks.

Lemma 2.27 *The functor i preserves injectives and we have an isomorphism $R(f_* \circ i) \cong Rf_* \circ Ri$.*

Proof. Since i^\sharp is exact (see [24, Thm. 3.2.1 (ii)]) the functor i preserves injectives. This implies the assertion. \square

2.3.5 Let $\tau := (U_i \rightarrow U)_{i \in I} \in \mathbf{cov}_{\mathbf{G}}(U \rightarrow G)$ be a covering family of $(U \rightarrow G) \in \mathbf{G}$. For a presheaf $F \in \mathbf{Pr}_{\mathbf{Ab}}\mathbf{G}$ we form the Čech complex $\check{C}^*(\tau, F)$. Its p th group is

$$\check{C}^p(\tau, F) := \prod_{(i_0, \dots, i_p) \in I^{p+1}} F(U_{i_0} \times_U \cdots \times_U U_{i_p}),$$

and the differential is given by the usual formula.

Definition 2.28 (see 3.5.1, [24]) *A sheaf $F \in \mathbf{Sh}_{\mathbf{Ab}}\mathbf{G}$ is called flabby if for all $(U \rightarrow G) \in \mathbf{G}$ and all $\tau \in \mathbf{cov}_{\mathbf{G}}(U \rightarrow G)$ we have $H^k(\check{C}(\tau, F)) \cong 0$ for all $k \geq 1$.*

2.3.6 Let $f : \mathbf{G} \rightarrow \mathbf{H}$ be a smooth map between smooth stacks.

Lemma 2.29 *The functor $f^* : \mathbf{Pr}_{\mathbf{Ab}}\mathbf{G} \rightarrow \mathbf{Pr}_{\mathbf{Ab}}\mathbf{H}$ preserves flabby sheaves.*

Proof. We have the functor $f_\sharp : \mathbf{G} \rightarrow \mathbf{H}$ given by $f_\sharp(V \rightarrow G) := (V \rightarrow G \rightarrow H)$ (see 2.2.7). By Lemma 2.7 we know that f^* preserves sheaves.

Let $(U \rightarrow G) \in \mathbf{G}$ and $\tau := (U_i \rightarrow U) \in \mathbf{cov}_{\mathbf{G}}(U)$. Observe that $f_\sharp\tau := (f_\sharp(U_i) \rightarrow f_\sharp(U))$ is a covering family of $f_\sharp U$ in \mathbf{H} .

Let $F \in \mathbf{Sh}_{\mathbf{Ab}}\mathbf{H}$. By Lemma 2.8 we have $f^*F(U) \cong F(f_\sharp U)$. We therefore have an isomorphism of complexes

$$\check{C}(\tau, f^*F) \cong \check{C}(f_\sharp\tau, F).$$

If F is in addition flabby, then the cohomology groups of the right-hand side in degree ≥ 1 vanish. \square

2.3.7 Let $f : G \rightarrow H$ be a representable map between smooth stacks.

Lemma 2.30 *If $F \in \mathbf{Sh}_{\mathbf{Ab}}\mathbf{G}$ is flabby, then F is $(f_* \circ i)$ -acyclic.*

Proof. Let $F \in \mathbf{Sh}_{\mathbf{Ab}}\mathbf{G}$ be flabby. We must show that $R^k(f_* \circ i)(F) \cong 0$ for $k \geq 1$. By Lemma 2.11 the functor f_* is exact so that $R^k(f_* \circ i)(F) \cong f_* \circ R^k i(F)$. Since F is injective it is flabby. Since flabby sheaves are i -acyclic by [24, Corollary 3.5.3] we get $R^k i(F) \cong 0$. \square

2.3.8 Let G be a smooth stack, $F \in D^+(\mathbf{Sh}_{\mathbf{Ab}}\mathbf{G})$, and $p : G \rightarrow *$ the canonical morphism. Then we have the object $Rp_* \circ Ri \in D^+(\mathbf{Pr}_{\mathbf{Ab}}\mathbf{Site}(*))$. Let $\mathbf{ev} : \mathbf{Pr}_{\mathbf{Ab}}\mathbf{Site}(*)) \rightarrow \mathbf{Ab}$ be the evaluation at the object $(* \rightarrow *) \in \mathbf{Site}(*))$. This functor is exact.

Definition 2.31 *We define the cohomology of $F \in D^+(\mathbf{Sh}_{\mathbf{Ab}}\mathbf{G})$ as*

$$h(G; F) := \mathbf{ev} \circ Rp_* \circ Ri(F) \in D^+(\mathbf{Ab})$$

Furthermore we set $H^(G; F) := H^*h(G; F)$.*

In particular, for an abelian group Z we have the constant presheaf Z_G with value Z .

Definition 2.32 *We define the cohomology of the smooth stack G with coefficients in Z by*

$$H^*(G; Z) := H^*(G; i^\sharp Z_G).$$

2.4 Simplicial models

2.4.1 For a morphism $f : G \rightarrow X$ between smooth stacks we defined a functor $f_* : \text{Pr } \mathbf{G} \rightarrow \text{Pr } \mathbf{X}$ (see Definition 2.4). We are in particular interested in its derived version $Rf_* \circ Ri : D^+(\text{Sh}_{\text{Ab}} \mathbf{G}) \rightarrow D^+(\text{Pr}_{\text{Ab}} \mathbf{X})$. The definitions of f_* in terms of a limit, and of Rf_* using injective resolutions are very useful for the study of the functorial properties of f_* . For explicit calculations we would like to work with more concrete objects. In the present subsection we associate to a flabby sheaf $F \in \text{Sh}_{\text{Ab}} \mathbf{G}$ an explicit complex of presheaves $C_A^+(F) \in C^+(\text{Pr}_{\text{Ab}} \mathbf{X})$ which represents $Rf_* \circ i(F) \in D^+(\text{Pr}_{\text{Ab}} \mathbf{X})$ (see Lemma 2.41). It looks like presheaf of Čech complexes and depends on the choice of a surjective smooth and representable map $A \rightarrow G$ such that $A \rightarrow G \rightarrow X$ is also representable (e.g. an atlas of G).

In the present paper we consider three applications of this construction. The first is the derived version of Lemma 2.16 which says that pull-back and push-forward in certain cartesian diagrams commute (see Lemma 2.43). In the second application we use the complex C_A^+ in order to get a de Rham model of the derived push-forward of the constant sheaf with value \mathbb{R} on \mathbf{G} (see equation (3.3)). Finally we use this construction in Lemma 3.8 in order to calculate the cohomology of the gerbe $[*/S^1]$ explicitly.

2.4.2 Let G be a smooth stack and $(A \rightarrow G), (B \rightarrow G) \in \mathbf{G}$.

Lemma 2.33 *The fibre product in stacks*

$$\begin{array}{ccccc}
 & & A \times_G B & & \\
 & \swarrow & \downarrow & \searrow & \\
 A & \overset{\dots\dots\dots}{\longleftarrow} & & \overset{\dots\dots\dots}{\longleftarrow} & B \\
 & \swarrow & \downarrow & \searrow & \\
 & & G & &
 \end{array}$$

$\begin{array}{ccc} \text{---} u \text{---} & & \text{---} \text{id} \text{---} \\ \text{---} & & \text{---} \end{array}$

is the categorial product $(A \rightarrow G) \times_{\mathbf{G}} (B \rightarrow G)$.

Proof. The fibre product $(H \rightarrow G), (L \rightarrow G) \mapsto H \times_G L$ of stacks $H, L \in \mathcal{C}$ over G is the two-categorical fibre product in the two-category \mathcal{C}/G of stacks over G . Let $\mathcal{C}_0 \subset \mathcal{C}$ be the full subcategory of stacks which are equivalent to smooth manifolds, i.e. the essential image of the Yoneda embedding $\text{Mf}^\infty \rightarrow \mathcal{C}$. We define the one-category $\overline{\mathcal{C}_0/G}$ by identifying two-isomorphic morphisms and observe that the canonical functor $\mathcal{C}_0/G \rightarrow \overline{\mathcal{C}_0/G}$ is an equivalence. Under this equivalence the restriction of the fibre product to \mathcal{C}_0 becomes the one-categorical product. This implies the result since the natural functor

$$\mathbf{G} \rightarrow \mathcal{C}_0/G \rightarrow \overline{\mathcal{C}_0/G}$$

is an equivalence of categories. □

2.4.3 Let $f : G \rightarrow X$ be a map of smooth stacks. Let further A be a smooth stack and $A \rightarrow G$ be a representable, surjective and smooth map such that the composition $A \rightarrow G \rightarrow X$ is also representable. An atlas of G would have these properties, but in applications we will need this more general situation where A is not necessarily equivalent to a manifold. Let $(U \rightarrow X) \in \mathbf{X}$ and form the following diagram of cartesian squares:

$$\begin{array}{ccc}
 A_U & \longrightarrow & A \\
 \downarrow & \nearrow j^U & \downarrow \\
 G_U & \longrightarrow & G \\
 \downarrow & \nearrow & \downarrow \\
 U & \longrightarrow & X
 \end{array} \quad . \quad (2.34)$$

Since smoothness is preserved by pull-back the horizontal maps are smooth. Since surjectivity is also preserved by pull-back the two upper vertical maps are surjective and smooth. Since $A \rightarrow X$ is representable, the stack $A_U \cong U \times_X A$ is equivalent to a manifold.

2.4.4 Note that $(A_U \rightarrow G_U) \in \mathbf{G}_U$. In view of Lemma 2.33 we can take powers of A_U in \mathbf{G}_U . Using these powers we form a simplicial object $A_U \in \mathbf{G}_U$. Its n -th object is given by

$$\underbrace{A_U \times_{G_U} \cdots \times_{G_U} A_U}_{n+1 \text{ factors}} \rightarrow G_U .$$

We let $j_!^U A_U \in \mathbf{G}$ denote the simplicial object in \mathbf{G} with n th object $(A_U^n \rightarrow G_U \xrightarrow{j^U} G)$. If $V \rightarrow U$ is a morphism in \mathbf{X} , then we obtain an induced morphism of simplicial objects $j_!^V A_V \rightarrow j_!^U A_U$ in \mathbf{G} .

2.4.5 If $F \in \text{Pr } \mathbf{G}$, then we consider the cosimplicial object $U \mapsto F(j_!^U A_U)$ in $\text{Pr } \mathbf{X}$. For a morphism $V \rightarrow U$ in \mathbf{G} the structure map $F(j_!^U A_U) \rightarrow F(j_!^V A_V)$ is induced by the morphism of simplicial objects $j_!^V A_V \rightarrow j_!^U A_U$ in \mathbf{G} .

Definition 2.35 For a presheaf of abelian groups $F \in \text{Pr}_{\text{Ab}} \mathbf{G}$ let

$$C_A(F) \in C^+(\text{Pr}_{\text{Ab}} \mathbf{X})$$

denote the chain complex of presheaves associated to the cosimplicial presheaf of abelian groups $U \mapsto F(j_!^U A_U)$. Its differential will be denoted by δ .


2.4.6 Let $F \in \text{Pr}_{\text{Ab}} \mathbf{G}$.

Lemma 2.36 We have a natural transformation $\psi : f_* F \rightarrow H^0 C_A(F)$ which is an isomorphism if F is a sheaf.

Proof. Let $(U \rightarrow X) \in \mathbf{X}$. We recall definition of the push-forward 2.1.5:

$$f_* F(U) = \lim_{(V \rightarrow G) \in \mathbf{G}/U} F(V) .$$

Observe that



belongs to \mathbf{G}/U so that we have an evaluation


$$\begin{array}{ccc} f_* F(U) & \xrightarrow{\text{evaluation}} & C_A^0(F)(U) \\ & \searrow \psi & \nearrow \\ & H^0 C_A(F)(U) & \end{array}$$

with a canonical factorization ψ by the definition of $H^0 C_A(F)(U)$ as a kernel.


Assume now that F is a sheaf. Then we must show that ψ is an isomorphism. Let

$$\begin{array}{ccc} V & \longrightarrow & G \\ \downarrow & \nearrow & \downarrow \\ U & \longrightarrow & X \end{array} \tag{2.37}$$

be in \mathbf{G}/U . Then we have a canonical factorization



Using the induced map $V \rightarrow G_U$ we form the diagram



We consider the composition $(V \times_{G_U} A_U \rightarrow A_U \rightarrow G)$ as an object in \mathbf{G} . Since $A_U \rightarrow G_U$ is smooth and surjective the map $(V \times_{G_U} A_U \rightarrow V)$ is a covering of V in \mathbf{G} (it is here where we use the big site). For a sheaf F we have

$$F(V) \cong \lim \left(F(A_U \times_{G_U} V) \rightrightarrows F((A_U \times_{G_U} V) \times_V (A_U \times_{G_U} V)) \right).$$

We further have

$$(A_U \times_{G_U} V) \times_V (A_U \times_{G_U} V) \cong A_U \times_{G_U} A_U \times_{G_U} V$$

and a diagram

$$\begin{array}{ccc}
 F(A_U \times_{G_U} V) & \rightrightarrows & F(A_U \times_{G_U} A_U \times_{G_U} V) \\
 \uparrow & & \uparrow \\
 F(j_!^U A_U^0) & \rightrightarrows & F(j_!^U A_U^1)
 \end{array}$$

(recall that $A_U^0 = A_U$ and $A_U^1 = A_U \times_{G_U} A_U$) induced by the projection along V . Since $H^0 C_A^*(F)(U)$ is the limit of the lower horizontal part the left vertical map induces a map $H^0 C_A^*(F)(U) \rightarrow F(V)$. Since this construction is natural in the object (2.37) of \mathbf{G}/U we obtain finally a map $H^0 C_A^*(F)(U) \rightarrow f_* F(U)$ which is the inverse to ψ . \square

2.4.7

Lemma 2.38 *If $F \in \text{Pr}_{\text{Ab}} \mathbf{G}$ is injective, then $H^i C_A^*(F) = 0$ for $i \geq 1$.*

Proof. We follow the ideas of the last part of the proof of [24, Thm. 2.2.3]. Let $(U \rightarrow X) \in \mathbf{X}$ and \mathcal{A}_U denote the simplicial presheaf of sets represented by $j_!^U A_U$. Furthermore, let $\mathbb{Z}_{\mathcal{A}_U}$ be the (non-positively graded) complex of free abelian presheaves generated by \mathcal{A}_U . Then for any presheaf $F \in \text{Pr}_{\text{Ab}} \mathbf{G}$ we have

$$C_A^*(F)(U) \cong \text{Hom}_{\text{Pr}_{\text{Ab}} \mathbf{G}}(\mathbb{Z}_{\mathcal{A}_U}, F).$$

Since F is injective $\text{Hom}_{\text{Pr}_{\text{Ab}} \mathbf{G}}(\dots, F)$ is an exact functor. Hence it suffices to show that $H^i(\mathbb{Z}_{\mathcal{A}_U}) = 0$ for $i \leq -1$. For $(V \rightarrow G) \in \mathbf{G}$ the complex $\mathbb{Z}_{\mathcal{A}_U}(V)$ is the complex associated to the linearization of the simplicial set $\text{Hom}_{\mathbf{G}}(V, j_!^U A_U)$. We now rewrite

$$A_U \times_{G_U} \cdots \times_{G_U} A_U \cong (A \times_G \cdots \times_G A) \times_X U \cong (A \times_G \cdots \times_G A) \times_G G_U. \quad (2.39)$$

We consider V , $(A \times_G \cdots \times_G A)$ and G_U with their canonical maps to G as objects of the two-category \mathcal{C}/G of stacks over G . The first object is a manifold and therefore does not have non-trivial two-automorphisms.

Since the maps $(A \times_G \cdots \times_G A) \rightarrow G$ and $G_U \rightarrow G$ are representable these objects of \mathcal{C}/G also do not have non-trivial two-automorphisms. By the same reasoning as in the proof of Lemma 2.33 we can interpret the fibre product (2.39) as a one-categorical product. We get

$$\begin{aligned} \mathrm{Hom}_{\mathbf{G}}(V, j_!^U A_U) &= \mathrm{Hom}_{\mathcal{C}/G}(V, (A \times_G \cdots \times_G A) \times_G G_U) \\ &\cong (\mathrm{Hom}_{\mathcal{C}/G}(V, A) \times \cdots \times \mathrm{Hom}_{\mathcal{C}/G}(V, A)) \times \mathrm{Hom}_{\mathcal{C}/G}(V, G_U) \\ &\cong \mathrm{Hom}_{\mathcal{C}/G}(V, A) \times \mathrm{Hom}_{\mathcal{C}/G}(V, G_U) \end{aligned}$$

For any set S , if we take the simplicial set S^\cdot of the powers of S , the complex associated to the linearization \mathbb{Z}_{S^\cdot} is exact in degrees ≤ -1 . Therefore the complex $\mathbb{Z}_{\mathrm{Hom}_{\mathcal{C}/G}(V, A)^\cdot}$ is exact in degree ≤ -1 . Since the tensor product with the free abelian group $\mathbb{Z}_{\mathrm{Hom}_{\mathcal{C}/G}(V, G_U)^\cdot}$ is an exact functor the complex

$$\mathbb{Z}_{\mathrm{Hom}_{\mathbf{G}}(V, j_!^U A_U)^\cdot} \cong \mathbb{Z}_{\mathrm{Hom}_{\mathcal{C}/G}(V, A)^\cdot} \otimes \mathbb{Z}_{\mathrm{Hom}_{\mathcal{C}/G}(V, G_U)^\cdot}$$

is exact in degree ≤ -1 , too. \square

2.4.8 Since exactness of complexes of presheaves is defined objectwise the functors $C_A^p : \mathrm{Pr}_{\mathrm{Ab}} \mathbf{G} \rightarrow \mathrm{Pr}_{\mathrm{Ab}} \mathbf{X}$ are exact for all $p \geq 0$. Composing with the total complex construction we extend the functor C_A^\cdot to a functor between the categories of lower bounded complexes $C_A : C^+(\mathrm{Pr}_{\mathrm{Ab}} \mathbf{G}) \rightarrow C^+(\mathrm{Pr}_{\mathrm{Ab}} \mathbf{X})$ (in order to distinguish this from the double complex we drop the \cdot at the symbol C_A). Since this functor is level-wise exact it descends to a functor $C_A : D^+(\mathrm{Pr}_{\mathrm{Ab}} \mathbf{G}) \rightarrow D^+(\mathrm{Pr}_{\mathrm{Ab}} \mathbf{X})$ between the lower-bounded derived categories.

2.4.9 Assume that F is a presheaf of associative algebras on \mathbf{G} . Then $C_A^\cdot(F)$ is a presheaf of DG -algebras in the following natural way. Pick

$$\alpha \in F(j_!^U A_U^p) \cong C_A^p(F)(U), \quad \beta \in F(j_!^U A_U^q) \cong C_A^q(F)(U).$$

We have natural maps $u : j_!^U A_U^{p+q} \rightarrow j_!^U A_U^p$ and $v : j_!^U A_U^{p+q} \rightarrow j_!^U A_U^q$ in \mathbf{G} projecting onto the first $p+1$ or last $q+1$ factors, respectively. Then we define $\alpha \cdot \beta \in F(j_!^U A_U^{p+q}) \cong C_A^{p+q}(F)(U)$ by $u^* \alpha \cdot v^* \beta$. One easily checks that $\delta(\alpha \cdot \beta) = \delta \alpha + (-1)^p \alpha \cdot \delta \beta$.

2.4.10 If F^\cdot is a presheaf of commutative DG -algebras, then $C_A(F^\cdot)$ is a presheaf of associative DG -algebras central over the presheaf of commutative DG -algebras $(\ker(\delta) : C_A^0(F^\cdot) \rightarrow C_A^1(F^\cdot))$.

2.4.11 By Lemma 2.36 we have a map

$$\psi : f_* F \rightarrow H^0 C_A(F)$$

which is an isomorphism if F is a sheaf.

Lemma 2.40 *For all $F \in \mathrm{Pr}_{\mathrm{Ab}} \mathbf{G}$ we have a natural isomorphism $RH^0 C_A(F) \cong C_A(F)$ in $D^+(\mathrm{Pr}_{\mathrm{Ab}} \mathbf{X})$.*

Proof. Let $F \rightarrow I$ be an injective resolution. Then we have $RH^0 C_A(F) \cong H^0 C_A(I)$. By Lemma 2.38 the inclusion $H^0 C_A(I) \rightarrow C_A(I)$ is a quasi-isomorphism. Since C_A^\cdot is exact the quasi-isomorphism $F \rightarrow I$ induces a quasi-isomorphism $C_A(F) \cong C_A(I)$. \square

2.4.12 Recall that $i : \mathrm{Pr}_{\mathrm{Ab}} \mathbf{G} \rightarrow \mathrm{Sh}_{\mathrm{Ab}} \mathbf{G}$ is left exact and admits a right derived functor $Ri : D^+(\mathrm{Sh}_{\mathrm{Ab}} \mathbf{G}) \rightarrow D^+(\mathrm{Pr}_{\mathrm{Ab}} \mathbf{G})$, and that C_A descends to a functor between the lower bounded derived categories (see 2.4.8).

Lemma 2.41 *We have a natural isomorphism of functors*

$$C_A \circ Ri \cong Rf_* \circ Ri : D^+(\mathrm{Sh}_{\mathrm{Ab}} \mathbf{G}) \rightarrow D^+(\mathrm{Pr}_{\mathrm{Ab}} \mathbf{X}).$$

Proof. By Lemma 2.36 we have an isomorphism of functors $f_* \circ i \cong H^0 C_A \circ i$. Hence we have an isomorphism

$$Rf_* \circ Ri \stackrel{!}{\cong} R(f_* \circ i) \cong R(H^0 C_A \circ i) \stackrel{!}{\cong} RH^0 C_A \circ Ri \cong C_A \circ Ri,$$

where at the marked isomorphisms we use that i preserves injectives (compare Lemma 2.27). \square

2.4.13 Assume that we have a diagram in smooth stacks

$$\begin{array}{ccc} G & \xrightarrow{u} & H \\ \downarrow f & \nearrow & \downarrow g \\ X & \xrightarrow{v} & Y \end{array} , \quad (2.42)$$

where u and v are smooth. Note that u^* and v^* are exact (Lemma 2.7).

Lemma 2.43 (1) We have a natural transformation of functors $D^+(\mathrm{Pr}_{\mathrm{Ab}} \mathbf{H}) \rightarrow D^+(\mathrm{Pr}_{\mathrm{Ab}} \mathbf{X})$

$$v^* \circ Rg_* \rightarrow Rf_* \circ u^* .$$

(2) The induced transformation $D^+(\mathrm{Sh}_{\mathrm{Ab}} \mathbf{H}) \rightarrow D^+(\mathrm{Pr}_{\mathrm{Ab}} \mathbf{X})$

$$v^* \circ Rg_* \circ Ri \rightarrow Rf_* \circ u^* \circ Ri$$

is an isomorphism if (2.42) is cartesian.

Proof. The transformation (1) is induced by

$$v^* \circ Rg_* \cong R(v^* \circ g_*) \stackrel{2.16}{\cong} R(f_* \circ u^*) \rightarrow Rf_* \circ u^* .$$

In order to show the second part (2) we must show that

$$R(f_* \circ u^*) \circ Ri \rightarrow Rf_* \circ u^* \circ Ri$$

is an isomorphism. We calculate Ri using injective resolutions. Note that i preserves injectives. Hence in order to show that this map is an isomorphism it suffices to show that u^* maps injective sheaves to f_* -acyclic presheaves.

Note that u^* preserves sheaves (Lemma 2.7). We let $u_s^* : \mathrm{Sh}_{\mathrm{Ab}} \mathbf{H} \rightarrow \mathrm{Sh}_{\mathrm{Ab}} \mathbf{G}$ denote the restriction of u^* to sheaves. Let $F \in \mathrm{Sh}_{\mathrm{Ab}} \mathbf{H}$ be injective. Since injective sheaves are flabby, flabby sheaves are i -acyclic, and u^* preserves flabby sheaves (see Lemma 2.29) we have

$$Rf_* \circ i \circ u_s^*(F) \cong Rf_* \circ Ri \circ u_s^*(F) \stackrel{\text{Lemma 2.41}}{\cong} C_A \circ Ri \circ u_s^*(F) \cong C_A \circ u^* \circ i(F) .$$

We now show that the higher cohomology presheaves of $C_A \circ u^* \circ i(F)$ vanish. Let $(U \rightarrow X) \in \mathbf{X}$ and choose an atlas $B \rightarrow H$. Then we get the following extension of the diagram (2.34)

$$\begin{array}{ccccc} A_U & \longrightarrow & A & \longrightarrow & B \\ \downarrow & \nearrow & \downarrow & \nearrow & \downarrow \\ G_U & \xrightarrow{j^U} & G & \xrightarrow{u} & H \\ \downarrow & \nearrow & \downarrow & \nearrow & \downarrow g \\ U & \longrightarrow & X & \xrightarrow{v} & Y \end{array} \quad (2.44)$$

such that all squares are cartesian. The three upper vertical maps are smooth and surjective. The composition $A \rightarrow G \rightarrow X$ is representable. All horizontal maps are smooth. We have the simplicial object $(A_U \rightarrow G_U) \in \mathbf{G}_U$ and let $u_! j_!^U A_U \in \mathbf{H}$ be the induced simplicial object $A_U \rightarrow G_U \xrightarrow{j^U} G \xrightarrow{u} H$ in \mathbf{H} . Then we have by the definition 2.4.5 of C_A and the formula (2.8) for u^* that

$$C_A \circ u^* \circ i(F)(U) = F(u_! j_!^U A_U) .$$

We now observe the isomorphisms

$$\begin{aligned} A_U \times_{G_U} \cdots \times_{G_U} A_U &\cong (A \times_G \cdots \times_G A) \times_X U \\ &\cong (B \times_H \cdots \times_H B) \times_Y U \\ &\cong B_{v_1 U} \times_{H_{v_1 U}} \cdots \times_{H_{v_1 U}} B_{v_1 U} , \end{aligned}$$

where the notation is explained by the cartesian diagram

$$\begin{array}{ccc} H_{v_1 U} & \xrightarrow{k} & H \\ \downarrow & & \downarrow \\ U & \longrightarrow & Y \end{array} ,$$

and where $v_1 U := (U \rightarrow X \xrightarrow{v} Y) \in \mathbf{Y}$. We can thus identify the simplicial object $u_! j_! A_U$ with the similar simplicial object $k_! B_{v_1 U}$ in \mathbf{H} . In other words, we have an isomorphism of complexes

$$C_A \circ u^* \circ i(F)(U) \cong C_B \circ i(F)(v_1 U) .$$

Since $i(F)$ is an injective presheaf the right-hand side is exact by Lemma 2.38. \square

2.5 Comparison of big and small sites

2.5.1 Let X be a smooth stack and $(U \rightarrow X) \in \mathbf{X}$. A presheaf on X naturally induces a presheaf on the small site (U) of the manifold U consisting of the open subsets. This restriction functor will be used subsequently in order to reduce assertions in the sheaf theory over X to assertions in the ordinary sheaf theory on U . The goal of the present subsection is to study exactness properties of this restriction and its relation with the sheafification functors.

2.5.2 If U is a smooth manifold, then we let (U) denote the small site of U where covering families are coverings by families of open submanifolds. A presheaf with respect to the big site on U is in particular a presheaf with respect to (U) .

2.5.3 Let G be a smooth stack and $(U \rightarrow G) \in \mathbf{G}$. Then we have a functor $\nu_U : \text{Pr } \mathbf{G} \rightarrow \text{Pr}(U)$ which associates to the presheaf $F \in \text{Pr } \mathbf{G}$ the presheaf $\nu_U(F) \in \text{Pr}(U)$ obtained by restriction of structure. Since limits and colimits in presheaves are defined objectwise the functor ν_U is exact.

2.5.4

Lemma 2.45 *The functor ν_U preserves sheaves and induces a functor $\nu_U^s : \text{Sh } \mathbf{G} \rightarrow \text{Sh}(U)$.*

Proof. An object $V \in (U)$ gives rise to an object $(V \rightarrow U \rightarrow G) \in \mathbf{G}$. Observe that covering families of objects of $V \in (U)$ are also covering families of $(V \rightarrow G) \in \mathbf{G}$. For open subsets $V_1, V_2 \subset V$ the fibre products $V_1 \times_V V_2$ in (U) and in \mathbf{G} coincide by the discussion in 2.1.3. Therefore the descent conditions on $\nu_U(F)$ to be a sheaf on (U) are part of the descent conditions for F to be a sheaf on \mathbf{G} . Hence the functor ν_U restricts to $\nu_U^s : \text{Sh } \mathbf{G} \rightarrow \text{Sh}(U)$. \square

2.5.5 Since limits of sheaves are defined objectwise the functor ν_U^s commutes with limits. The goal of the following discussion is to show that it also commutes with colimits.

Proposition 2.46 *The functor $\nu_U^s : \text{Sh } \mathbf{G} \rightarrow \text{Sh}(U)$ is exact.*

Proof. If F is a diagram of sheaves, then we have

$$\text{colim}^s(F) \cong i^\# \circ \text{colim} \circ i(F) ,$$

where colim^s is the colimit of sheaves. Note that $\nu_U \circ i \cong i \circ \nu_U^s$ and $\nu_U \circ \text{colim} \cong \text{colim} \circ \nu_U$. In order to show that ν_U^s commutes with colim^s it remains to show the following Lemma.

2.5.6

Lemma 2.47 *We have*

$$i^\# \circ \nu_U \cong \nu_U^s \circ i^\# : \Pr \mathbf{G} \rightarrow \mathbf{Sh}(U) .$$

Proof. For the moment it is useful to indicate by a subscript (e.g. $i_{\mathbf{G}}$ or $i_{(U)}$) the site for which the functors are considered. Following the discussion in [24, Section 3.1] we introduce an explicit construction of the sheafification functor. Consider the site \mathbf{G} . We define the functor $P_{\mathbf{G}} : \Pr \mathbf{G} \rightarrow \Pr \mathbf{G}$ as follows. Let $(V \rightarrow G) \in \mathbf{G}$. Then we have the category of covering families $\mathbf{cov}_{\mathbf{G}}(V)$ whose morphisms are refinements. For $\tau := (V_i \rightarrow V) \in \mathbf{cov}_{\mathbf{G}}(V)$ we define $H^0(F)(\tau)$ by the equalizer diagram

$$H^0(F)(\tau) \rightarrow \prod_i F(V_i) \rightrightarrows \prod_{i,j} F(V_i \times_V V_j) .$$

We get a diagram $\tau \rightarrow H^0(F)(\tau)$ in $\mathbf{Sets}^{\mathbf{cov}_{\mathbf{G}}(V)}$ and define

$$P_{\mathbf{G}}(F)(V) := \operatorname{colim}_{\tau \in \mathbf{cov}_{\mathbf{G}}(V)} H^0(F)(\tau) .$$

Then we have

$$i_{\mathbf{G}} \circ i_{\mathbf{G}}^\# := P_{\mathbf{G}} \circ P_{\mathbf{G}} : \Pr \mathbf{G} \rightarrow \Pr \mathbf{G} .$$

In a similar manner we define a functor $P_{(U)} : \Pr(U) \rightarrow \Pr(U)$ and get

$$i_{(U)} \circ i_{(U)}^\# := P_{(U)} \circ P_{(U)} : \Pr(U) \rightarrow \Pr(U) .$$

In order to show the Lemma it suffices to show that

$$P_{(U)} \circ \nu_U \cong \nu_U \circ P_{\mathbf{G}} .$$

Let $V \subset U$ be open and consider the induced $(V \rightarrow G) \in \mathbf{G}$. Then we have a functor

$$a : \mathbf{cov}_{(U)}(V) \rightarrow \mathbf{cov}_{\mathbf{G}}(V) . \quad (2.48)$$

If $\tau \in \mathbf{cov}_{(U)}(V)$, then we have an isomorphism

$$H^0(F)(a(\tau)) \cong H^0(\nu_U(F))(\tau) .$$

We therefore have an induced map of colimits

$$P_{(U)} \circ \nu_U(F)(V) \rightarrow \nu_U \circ P_{\mathbf{G}}(F)(V) .$$

This map is in fact an isomorphism since we will show below that (2.48) defines a cofinal subfamily. Let $\sigma := (U_i \rightarrow V)_{i \in I} \in \mathbf{cov}_{\mathbf{G}}(V)$. Since the maps $U_i \rightarrow V$ are submersions they admit local sections. Hence there exists a covering $\tau := (V_j \rightarrow V)_{j \in J} \in \mathbf{cov}_{(U)}$, a map $r : J \rightarrow I$ and a family of sections $s_j : V_j \rightarrow U_{r(j)}$ such that

$$\begin{array}{ccc} & U_{r(j)} & \\ s_j \nearrow & & \searrow \\ V_j & \xrightarrow{\quad} & V \end{array}$$

commutes for all $j \in J$. This data defines a morphism $\sigma \rightarrow a(\tau)$ in $\mathbf{cov}_{\mathbf{G}}(V)$. □

This finishes the proof of Proposition 2.46. □

2.5.7 Recall the definition of a flabby sheaf 2.28.

Lemma 2.49 *The functor $\nu_U^s : \mathbf{Sh}_{\text{Ab}} \mathbf{G} \rightarrow \mathbf{Sh}_{\text{Ab}}(U)$ preserves flabby sheaves.*

Proof. Let $V \subset U$ be an open subset and $\tau \in \text{cov}_{(U)}(V)$. Let $a : \text{cov}_{(U)}(V) \rightarrow \text{cov}_{\mathbf{G}}(V)$ be as in (2.48). We have an natural isomorphism $\check{C}(\tau, \nu_U^s(F)) \cong \check{C}(a(\tau), F)$. If $F \in \mathbf{Sh}_{\text{Ab}} \mathbf{G}$ is flabby, then for $k \geq 1$ we have $H^k \check{C}(\tau, \nu_U^s(F)) \cong H^k(\check{C}(a(\tau), F)) \cong 0$. \square

2.5.8 Since $\nu_U : \text{Pr } \mathbf{G} \rightarrow \text{Pr}(U)$ is exact it descends to a functor $\nu_U : D^+(\text{Pr}_{\text{Ab}} \mathbf{G}) \rightarrow D^+(\text{Pr}_{\text{Ab}}(U))$ between the lower bounded derived categories. Since $\nu_U^s : \mathbf{Sh} \mathbf{G} \rightarrow \mathbf{Sh}(U)$ is exact, it descends to a functor $\nu_U^s : D^+(\mathbf{Sh}_{\text{Ab}} \mathbf{X}) \rightarrow D^+(\mathbf{Sh}_{\text{Ab}}(U))$.

2.5.9 Using the techniques above we show the following result which will be useful later. Let $f : G \rightarrow H$ be a smooth map between smooth stacks. Note that $f^* : \text{Pr}_{\text{Ab}} \mathbf{H} \rightarrow \text{Pr}_{\text{Ab}} \mathbf{G}$ is exact.

Lemma 2.50 (1) *We have an isomorphism of functors*

$$f^* \circ i_{\mathbf{H}} \circ i_{\mathbf{H}}^{\sharp} \cong i_{\mathbf{G}} \circ i_{\mathbf{G}}^{\sharp} \circ f^* : \text{Pr}_{\text{Ab}} \mathbf{H} \rightarrow \text{Pr}_{\text{Ab}} \mathbf{G} .$$

(2) *We have an isomorphism of functors*

$$f^* \circ Ri_{\mathbf{H}} \circ i_{\mathbf{H}}^{\sharp} \cong Ri_{\mathbf{G}} \circ i_{\mathbf{G}}^{\sharp} \circ f^* : D^+(\text{Pr}_{\text{Ab}} \mathbf{H}) \rightarrow D^+(\text{Pr}_{\text{Ab}} \mathbf{G}) .$$

Proof. Let $(U \rightarrow G) \in \mathbf{G}$ and $f_!U := (U \rightarrow G \xrightarrow{f} H) \in \mathbf{H}$. We calculate for $F \in \text{Pr}_{\text{Ab}} \mathbf{H}$ that on the one hand

$$\begin{aligned} (f^* \circ i_{\mathbf{H}} \circ i_{\mathbf{H}}^{\sharp} F)(U) &\stackrel{(2.7)}{\cong} (i_{\mathbf{H}} \circ i_{\mathbf{H}}^{\sharp} F)(f_!U) \\ &\cong (\nu_{f_!U} \circ i_{\mathbf{H}} \circ i_{\mathbf{H}}^{\sharp} F)(U) \\ &\stackrel{2.45, 2.47}{\cong} (i_{(U)} \circ i_{(U)}^{\sharp} \circ \nu_{f_!U} F)(U) . \end{aligned}$$

On the other hand we have

$$\begin{aligned} (i_{\mathbf{G}} \circ i_{\mathbf{G}}^{\sharp} \circ f^* F)(U) &\cong (\nu_U \circ i_{\mathbf{G}} \circ i_{\mathbf{G}}^{\sharp} \circ f^* F)(U) \\ &\stackrel{2.45, 2.47}{\cong} (i_{(U)} \circ i_{(U)}^{\sharp} \circ \nu_U \circ f^* F)(U) . \end{aligned}$$

Finally we use the fact that $\nu_U \circ f^* F \cong \nu_{f_!U} F$. Indeed, for $V \subset U$ we have

$$\nu_U \circ f^* F(V) \cong F(f_!V) \cong \nu_{f_!U} F(V) .$$

The combination of these isomorphisms gives the first assertion.

Since f^* preserves sheaves we can consider the restriction $f_s^* : \mathbf{Sh}_{\text{Ab}} \mathbf{H} \rightarrow \mathbf{Sh}_{\text{Ab}} \mathbf{G}$ of f^* . Using the first part of the Lemma and the isomorphism $i^{\sharp} \circ i \cong \text{id}$ we get

$$f_s^* \circ i_{\mathbf{H}}^{\sharp} \cong i_{\mathbf{G}}^{\sharp} \circ i_{\mathbf{G}} \circ f_s^* \circ i_{\mathbf{H}}^{\sharp} \cong i_{\mathbf{G}}^{\sharp} \circ f^* \circ i_{\mathbf{H}} \circ i_{\mathbf{H}}^{\sharp} \stackrel{(1)}{\cong} i_{\mathbf{G}}^{\sharp} \circ i_{\mathbf{G}} \circ i_{\mathbf{G}}^{\sharp} \circ f^* \cong i_{\mathbf{G}}^{\sharp} \circ f^* . \quad (2.51)$$

Note that f_s^* is an exact functor. In order to see that it is right exact we use that f_* preserves sheaves we consider its restriction f_*^s to sheaves. For $X \in \mathbf{Sh} \mathbf{H}$ and $Y \in \mathbf{Sh} \mathbf{G}$ we get a natural isomorphism

$$\begin{aligned} \text{Hom}_{\mathbf{Sh} \mathbf{G}}(f_s^* X, Y) &\cong \text{Hom}_{\mathbf{Sh} \mathbf{G}}(i_{\mathbf{G}}^{\sharp} \circ i_{\mathbf{G}} f_s^* X, Y) \\ &\cong \text{Hom}_{\text{Pr } \mathbf{G}}(i_{\mathbf{G}} f_s^* X, i_{\mathbf{G}} Y) \\ &\cong \text{Hom}_{\text{Pr } \mathbf{G}}(f^* \circ i_{\mathbf{H}} X, i_{\mathbf{G}} Y) \\ &\cong \text{Hom}_{\text{Pr } \mathbf{H}}(i_{\mathbf{H}} X, f_* \circ i_{\mathbf{G}} Y) \\ &\cong \text{Hom}_{\text{Pr } \mathbf{H}}(i_{\mathbf{H}} X, i_{\mathbf{H}} \circ f_*^s Y) \\ &\cong \text{Hom}_{\mathbf{Sh} \mathbf{H}}(X, f_*^s Y) \end{aligned}$$

Therefore f_s^* is a left adjoint and therefore right exact. We now write

$$f_s^* \cong i_{\mathbf{G}}^\sharp \circ i_{\mathbf{G}} \circ f_s^* \cong i_{\mathbf{G}}^\sharp \circ f^* \circ i_{\mathbf{H}} .$$

Since $i_{\mathbf{G}}^\sharp$ is exact, and f^* and $i_{\mathbf{H}}$ are left exact (since they are right adjoints, see Lemma 2.25 for a left adjoint of f^*) we conclude that f_s^* is left exact, too

Using the that f_s^* preserves flabby sheaves (Lemma 2.29) and that it is an exact functor we get

$$f^* \circ Ri_{\mathbf{H}} \cong R(f^* \circ i_{\mathbf{H}}) \cong R(i_{\mathbf{G}} \circ f_s^*) \cong Ri_{\mathbf{G}} \circ f_s^* .$$

Combining this with (2.51) we get the desired isomorphism

$$f^* \circ Ri_{\mathbf{H}} \circ i_{\mathbf{H}}^\sharp \cong Ri_{\mathbf{G}} \circ f_s^* \circ i_{\mathbf{H}}^\sharp \cong Ri_{\mathbf{G}} \circ i_{\mathbf{G}}^\sharp \circ f^* .$$

□

3 The de Rham complex

3.1 The de Rham complex is a flabby resolution

3.1.1 We want to apply Lemma 2.41 to the sheafification $i_{\mathbf{G}}^\sharp \mathbb{R}_{\mathbf{G}}$ of the constant presheaf with value \mathbb{R} on \mathbf{G} . In particular, we must calculate $Ri(i_{\mathbf{G}}^\sharp \mathbb{R}_{\mathbf{G}})$. This can be done by applying i to a flabby resolution of $i_{\mathbf{G}}^\sharp \mathbb{R}_{\mathbf{G}}$. In the present subsection we introduce the de Rham complex G and show that it is a flabby resolution of $i_{\mathbf{G}}^\sharp \mathbb{R}_{\mathbf{G}}$.

The de Rham complex of G is build from the de Rham complexes of the manifolds U for all $(U \rightarrow G) \in \mathbf{G}$. For each U equipped with the topology of the small site it is well known that the de Rham complex resolves the constant sheaf with value \mathbb{R} and is flabby. Our task here is to extend these properties to the stack G and the big site.

3.1.2 Let G be a smooth stack and fix an integer $p \geq 0$. We define the presheaf $\Omega^p(G)$ by

$$\Omega^p(G)(U) := \Omega^p(U) .$$

If $\phi : U \rightarrow V$ is a morphism in \mathbf{G} , then $\Omega^p(G)(\phi) := \phi^* : \Omega^p(V) \rightarrow \Omega^p(U)$. Since ϕ^* commutes with the de Rham differential we get a complex $(\Omega^*(G), d_{dR})$ of presheaves.

Lemma 3.1 *The presheaf $\Omega^p(G)$ is a sheaf and flabby.*

Proof. Let $(U \rightarrow G) \in \mathbf{G}$. Observe that $\nu_U(\Omega^p(G))$ is the presheaf of smooth sections of the vector bundle $\Lambda^p T^*U$. This is actually a sheaf. In order to show that $\Omega^p(G)$ is a sheaf it suffices to show that the unit $\Omega^p(G) \rightarrow i_{\mathbf{G}} \circ i_{\mathbf{G}}^\sharp(\Omega^p(G))$ of the adjoint pair $(i_{\mathbf{G}}^\sharp, i_{\mathbf{G}})$ is an isomorphism. This follows from the calculation

$$\begin{aligned} i_{\mathbf{G}} \circ i_{\mathbf{G}}^\sharp(\Omega^p(G))(U) &\cong \nu_U \circ i_{\mathbf{G}} \circ i_{\mathbf{G}}^\sharp(\Omega^p(G))(U) \\ &\stackrel{\text{Lemma 2.45}}{\cong} i_{(U)} \circ \nu_U^s \circ i_{\mathbf{G}}^\sharp(\Omega^p(G))(U) \\ &\stackrel{\text{Lemma 2.47}}{\cong} i_{(U)} \circ i_{(U)}^\sharp \circ \nu_U(\Omega^p(G))(U) \\ &\stackrel{\nu_U(\Omega^p(G)) \text{ is a sheaf}}{\cong} \nu_U(\Omega^p(G))(U) . \end{aligned}$$

A sheaf $F \in \mathbf{Sh}_{\text{Ab}}(U)$ on a paracompact space U is called soft if for all closed subsets $Z \subset U$ the restriction $\Gamma_U(F) \rightarrow \Gamma_Z(F)$ is surjective. For a soft sheaf we have $R^i \Gamma_U(F) \cong 0$ for all $i \geq 1$ (see [13, Ex. II 5]). It now follows from [24, Corollary 3.5.3] that a soft sheaf is flabby.

A sheaf of smooth sections of a smooth vector bundle on a smooth manifold is soft. In particular, $\nu_U^s(\Omega^p(G))$ is soft and therefore flabby.

In order to show that the sheaf $\Omega^p(G)$ is flabby it suffices by [24, Corollary 3.5.3] to show that $R^k i(\Omega^p(G)) \cong 0$ for $k \geq 1$. We calculate

$$\begin{aligned}
R^k i(\Omega^p(G))(U) &\cong \Gamma_U \circ R^k i(\Omega^p(G)) \\
&\stackrel{\Gamma_U \text{ exact}}{\cong} H^k(\Gamma_U \circ Ri(\Omega^p(G))) \\
&\stackrel{H^k \text{ objectwise}}{\cong} H^k(Ri(\Omega^p(G))(U)) \\
&\stackrel{\text{definition of } \nu_U}{\cong} H^k(\nu_U \circ Ri_{\mathbf{G}}(\Omega^p(G))(U)) \\
&\stackrel{\nu_U \text{ exact}}{\cong} R^k(\nu_U \circ i_{\mathbf{G}})(\Omega^p(G))(U) \\
&\stackrel{\text{Lemma 2.45}}{\cong} R^k(i_{(U)} \circ \nu_U^s)(\Omega^p(G))(U) \\
&\stackrel{\text{Lemma 2.49}}{\cong} R^k i_{(U)} \circ \nu_U^s(\Omega^p(G))(U) \\
&\stackrel{\nu_U^s(\Omega^p(G)) \text{ is flabby}}{\cong} 0 .
\end{aligned}$$

□

3.1.3 Let $\mathbb{R}_{\mathbf{G}}$ denote the constant presheaf on \mathbf{G} with value \mathbb{R} and $i^{\sharp} \mathbb{R}_{\mathbf{G}}$ its sheafification. We have a canonical map $\mathbb{R}_{\mathbf{G}} \rightarrow H^0(\Omega^{\cdot}(G))$. Since $H^0(\Omega^{\cdot}(G))$ is a sheaf we get an induced map $i^{\sharp} \mathbb{R}_{\mathbf{G}} \rightarrow H^0(\Omega^{\cdot}(G))$.

Lemma 3.2 *The map*

$$i^{\sharp} \mathbb{R}_{\mathbf{G}} \rightarrow \Omega^{\cdot}(G)$$

is a quasi-isomorphism.

Proof. Note that the cohomology sheaves $H_s^k(F^{\cdot}) \in \mathbf{Sh}_{\text{Ab}} \mathbf{G}$ of a complex F^{\cdot} of sheaves of abelian groups on \mathbf{G} are defined by $H_s^k(F^{\cdot}) := i_{\mathbf{G}}^{\sharp} \circ H^k \circ i_{\mathbf{G}}(F^{\cdot})$, where H^k takes the cohomology of a complex of presheaves objectwise. We calculate

$$\begin{aligned}
H_s^k(\Omega^{\cdot}(G))(U) &\stackrel{\text{definition of } H_s^k}{\cong} (i_{\mathbf{G}}^{\sharp} \circ H^k \circ i_{\mathbf{G}})(\Omega^{\cdot}(G))(U) \\
&\stackrel{\text{Lemma 2.45}}{\cong} (\nu_U^s \circ i_{\mathbf{G}}^{\sharp} \circ H^k \circ i_{\mathbf{G}})(\Omega^{\cdot}(G))(U) \\
&\stackrel{\text{Lemma 2.47}}{\cong} (i_{(U)}^{\sharp} \circ \nu_U \circ H^k \circ i_{\mathbf{G}})(\Omega^{\cdot}(G))(U) \\
&\stackrel{\nu_U \text{ is exact}}{\cong} (i_{(U)}^{\sharp} \circ H^k \circ \nu_U \circ i_{\mathbf{G}})(\Omega^{\cdot}(G))(U) \\
&\stackrel{\text{Lemma 2.45}}{\cong} (i_{(U)}^{\sharp} \circ H^k \circ i_{(U)} \circ \nu_U^s)(\Omega^{\cdot}(G))(U) \\
&\stackrel{\text{definition of } H_s^k}{\cong} (H_s^k \circ \nu_U^s)(\Omega^{\cdot}(G))(U) .
\end{aligned}$$

Since $\nu_U^s(\Omega^{\cdot}(G))$ is the de Rham complex of the manifold U its higher cohomology sheaves vanish by the Poincaré Lemma. This implies that $H^k(\Omega^{\cdot}(G)) \cong 0$ for $k \geq 1$.

Furthermore, it is well-known that $H^0(\nu_U(\Omega^{\cdot}(G))) \cong i_{(U)}^{\sharp} \mathbb{R}_{(U)}$. It follows from the observation

$$\nu_U^s \circ i_{\mathbf{G}}^{\sharp} \mathbb{R}_{\mathbf{G}} \cong i_{(U)}^{\sharp} \mathbb{R}_{(U)}$$

(proved by arguments similar as above) that $i_{\mathbf{G}}^{\sharp} \mathbb{R}_{\mathbf{G}} \cong H^0(\Omega^{\cdot}(G))$. □

3.1.4 By Lemmas 3.1 and 3.2 the complex $\Omega^\cdot(G)$ is a flabby resolution of $i^\sharp \mathbb{R}_G$. Therefore

$$Ri(i^\sharp \mathbb{R}_G) \cong i(\Omega^\cdot(G))$$

in $D^+(\text{Pr}_{\text{Ab}} \mathbf{G})$. By Lemma 2.41 we have the isomorphism

$$(Rf_* \circ Ri)(i^\sharp \mathbb{R}_G) \cong C_A(\Omega^\cdot(G)) \quad (3.3)$$

in $D^+(\text{Pr}_{\text{Ab}} \mathbf{X})$.

3.2 Calculation for $U(1)$ -gerbes

3.2.1 In this subsection we specialize the situation of 2.4.3 to the case where $f : G \rightarrow X$ is a gerbe with band $U(1)$ according to the definition in 1.3.14 over a manifold X . We thus can assume that $A \rightarrow X$ is an atlas obtained from a covering of X by open subsets such that the lift $s : A \rightarrow G$ is an atlas of G . The $U(1)$ -central extension of groupoids in manifolds $(A \times_G A \rightrightarrows A) \rightarrow (A \times_X A \rightrightarrows A)$ (we forget the structure maps to G for the moment) is the picture of a gerbe as presented in [11]. In order to compare the sheaf theoretic construction of the cohomology of G with the twisted de Rham complex we must choose some additional geometric structure on G , namely a connection in the sense of [11]. The comparison map will depend on this choice.

3.2.2 A connection on the gerbe $f : G \rightarrow X$ consists of a pair (α, β) , where $\alpha \in \Omega^1(A \times_G A)$ is a connection one-form on the $U(1)$ -bundle $A \times_G A \rightarrow A \times_X A$, and $\beta \in \Omega^2(A)$. Observe that $\Omega^2(A)$ and $\Omega^1(A \times_G A)$ are the first two spaces of the degree-two part of the graded commutative DG-algebra $C_A(\Omega^\cdot(G))(X) \cong \Omega^\cdot(G)(A)$ discussed in 2.4.9 and 2.4.5. To be a connection the pair (α, β) is required to satisfy the following two conditions:

- (1) $\delta\beta = d_{dR}\alpha$ (where δ is the Čech differential of the complex $\Omega^\cdot(A)$, and d_{dR} is the de Rham differential) and
- (2) $\delta\alpha = 0$.

Note that $\delta d_{dR}\beta = 0$ so that there is a unique $\lambda \in \Omega^3(X)$ which restricts to $d_{dR}\beta$. We have $d_{dR}\lambda = 0$, and the class $[\lambda] \in H^3(X; \mathbb{R})$ represents the image under $H^3(X; \mathbb{Z}) \rightarrow H^3(X; \mathbb{R})$ of the Dixmier-Douady class of the gerbe $G \rightarrow X$ (see [11] for this fact and the existence of connections).

3.2.3 Let us choose a connection (α, β) , and let $\lambda \in \Omega^3(X)$ be the associated closed three form. We consider $(\alpha, \beta) \in C_A(\Omega^\cdot(G))^2(X)$.

We consider the sheaf of complexes $\Omega^\cdot[[z]]_\lambda$ on \mathbf{X} which associates to $(U \xrightarrow{i} X) \in \mathbf{X}$ the complex

$$\Omega^\cdot(U)[[z]], \quad d_\lambda := d_{dR} + \lambda T,$$

where $T := \frac{d}{dz}$, z has degree two, and λ acts by right multiplication by $i^*\lambda$. In particular we have $d_\lambda z = i^*\lambda$. Note that $\Omega^\cdot[[z]]_\lambda$ is a sheaf of left Ω_X -DG-algebras.

3.2.4 Observe that $z \in \Omega^\cdot[[z]]_\lambda(X)$ is central. Let $L \in \text{Pr}_{\text{Ab}} \mathbf{X}$ be a presheaf of graded unital central Ω_X -algebras. A map of presheaves of graded unital central Ω_X -algebras $\phi : i\Omega^\cdot[[z]]_\lambda \rightarrow L$ determines a section $\phi(z) \in L(X)$. Vice versa, given a section $l \in L(X)$ of degree two, there is a unique map of presheaves of graded unital central Ω_X -algebras $\phi : i\Omega^\cdot[[z]]_\lambda \rightarrow L$ such that $\phi(z) = l$. For $(U \xrightarrow{i} X) \in \mathbf{X}$ the map $\phi_U : i\Omega^\cdot[[z]]_\lambda(U) \rightarrow L(U)$ is given by

$$\phi_U\left(\sum_{k \geq 0} \omega_k z^k\right) := \sum_{k \geq 0} \omega_k i^*(l)^k,$$

where $i^* : L(X) \rightarrow L(U)$ is determined by the presheaf structure of L .

If (L, d^L) is a presheaf of DG-algebras over Ω_X , then ϕ is a homomorphism of DG-algebras over Ω_X if and only if $d^L l = \lambda$.

3.2.5

Proposition 3.4 *We have an isomorphism*

$$\Omega[[z]]_\lambda \cong i^\# C_A(\Omega(G))$$

in $D^+(\mathbf{Sh}_{\text{Ab}} \mathbf{X})$.

Proof. $C_A(\Omega(G))$ is a presheaf of DG -algebras by the 2.4.10. Given $(U \rightarrow X) \in \mathbf{X}$ we have a natural projection $\pi : A_U^0 \rightarrow U$ (see 2.4.5 for the notation). It induces a homomorphism of DG -algebras $\Omega_X(U) \rightarrow (\ker(\delta) : \Omega(G)(A_U^0) \rightarrow \Omega(G)(A_U^1))$ and therefore on $\Omega(G)(A_U^1)$ the structure of an $\Omega_X(U)$ - DG -module (see 2.4.10). In this way $C_A(\Omega(G))$ becomes a sheaf of central Ω_X - DG -algebras.

By the discussion in 3.2.4 we can define a map of presheaves of central Ω_X -algebras

$$\tilde{\phi} : i\Omega[[z]]_\lambda \rightarrow C_A(\Omega(G))$$

such that $\tilde{\phi}(z) = (\alpha, \beta) \in C_A(\Omega(G))^2(X)$. Because of $d\tilde{\phi}(z) = d(\alpha, \beta) = \lambda$, the map $\tilde{\phi}$ is a map of presheaves of DG -algebras over Ω_X , hence in particular a map of presheaves of complexes.

We let

$$\phi : \Omega[[z]]_\lambda \xrightarrow{\sim} i^\# \circ i\Omega[[z]]_\lambda \xrightarrow{i^\# \tilde{\phi}} i^\# C_A(\Omega(G))$$

be the induced map, where the first isomorphism exists since $\Omega[[z]]_\lambda$ is a complex of sheaves.

3.2.6 It remains to show that ϕ is a quasi-isomorphism of complexes of sheaves. This can be shown locally. We can therefore assume that X is contractible. We then have a pull-back diagram

$$\begin{array}{ccc} G & \xrightarrow{q} & [* / S^1] \\ \downarrow f & & \downarrow g \\ X & \xrightarrow{p} & * \end{array} .$$

Since p is smooth, so is q . By Lemma 2.43 we have a canonical isomorphism

$$p^* \circ Rg_* \circ Ri \xrightarrow{\sim} Rf_* \circ q^* \circ Ri . \quad (3.5)$$

Applying (3.5) to $i^\# \mathbb{R}_{\text{Site}([* / S^1])}$ we obtain

$$p^* \circ Rg_* \circ Ri(i^\# \mathbb{R}_{\text{Site}([* / S^1])}) \xrightarrow{\sim} Rf_* \circ q^* \circ Ri(i^\# \mathbb{R}_{\text{Site}([* / S^1])}) \quad (3.6)$$

in $D^+(\mathbf{Pr}_{\text{Ab}} \mathbf{X})$. We now use (see 3.1.4) that

$$Ri(i^\# \mathbb{R}_{\text{Site}([* / S^1])}) \cong i(\Omega([* / S^1])) .$$

By the calculation of q^* in Lemma 2.7 and the definition of the de Rham complex we have

$$q^* \circ i(\Omega([* / S^1])) \cong i(\Omega(G)) .$$

Therefore in $D^+(\mathbf{Pr}_{\text{Ab}} \mathbf{G})$ we have

$$q^* \circ Ri(i^\# \mathbb{R}_{\text{Site}([* / S^1])}) \cong i(\Omega(G)) \stackrel{3.1.4}{\cong} Ri(i^\# \mathbb{R}_{\mathbf{G}}) .$$

It follows by 3.1.4 that

$$Rf_* \circ q^* \circ Ri(i^\# \mathbb{R}_{\text{Site}([* / S^1])}) \cong Rf_* \circ Ri(i^\# \mathbb{R}_{\mathbf{G}}) \cong C_A(\Omega(G)) . \quad (3.7)$$

3.2.7 We now must calculate the cohomology of the gerbe $[*/S^1]$ with real coefficients.

Lemma 3.8 *We have an isomorphism*

$$i^\sharp \circ Rg_* \circ Ri(i^\sharp \mathbb{R}_{\mathbf{Site}([*/S^1])}) \cong i^\sharp(\mathbb{R}[[z]]_{\mathbf{Site}(*)}),$$

where z has degree two.

Proof. We choose the atlas $A := * \rightarrow [*/S^1]$ and use the isomorphism

$$Rg_* \circ Ri(i^\sharp \mathbb{R}_{\mathbf{Site}([*/S^1])}) \cong C_A(\Omega^{\cdot}([*/S^1])) \in D^+(\mathbf{Pr}_{\mathbf{Ab}}\mathbf{Site}(*)).$$

Note that $\mathbf{Site}(*)$ is the category of smooth manifolds. Let U be a smooth manifold. We have

$$A_U \cong U \rightarrow U \times [*/S^1] \cong [*/S^1]_U$$

and

$$C_A(\Omega^{\cdot}([*/S^1]))(U) \cong \Omega^{\cdot}(A_U),$$

where

$$A_U^p = \underbrace{A_U \times_{[*/S^1]_U} \cdots \times_{[*/S^1]_U} A_U}_{p+1 \text{ factors}} \cong U \times \underbrace{(* \times_{[*/S^1]} \cdots \times_{[*/S^1]} *)}_{p+1 \text{ factors}} \cong U \times (S^1)^p.$$

The simplicial manifold

$$A_U \cong U \times (S^1)^{\cdot}$$

is the simplicial model of the space $U \times BS^1$, where BS^1 is the classifying space of the group S^1 . We can use the simplicial de Rham complex in order to calculate its cohomology. Note that $H^*(BS^1, \mathbb{R}) \cong \mathbb{R}[[z]]$ with z in degree two. Let us fix a form $\zeta \in (\Omega^{\cdot}((S^1)^{\cdot})_{tot}^2)$ which represents the generator z . Then we define a map

$$\mu_U : \Omega^{\cdot}(U)[[z]] \rightarrow \Omega^{\cdot}(U \times (S^1)^{\cdot})$$

by

$$\mu(\omega z^k) := \omega \wedge \zeta^k.$$

This map induces a quasi-isomorphism of complexes of abelian groups. The family of maps μ_U for varying U defines a quasi-isomorphism of complexes of presheaves $\mu : i\Omega^{\cdot}(*)[[z]] \rightarrow C_A(\Omega^{\cdot}([*/S^1]))$. It induces the quasi-isomorphism of complexes of sheaves

$$\Omega^{\cdot}(*)[[z]] \cong i^\sharp \circ i\Omega^{\cdot}(*)[[z]] \xrightarrow{i^\sharp \mu} i^\sharp C_A(\Omega^{\cdot}([*/S^1])).$$

Finally observe that the canonical map

$$i^\sharp \mathbb{R}[[z]]_{\mathbf{Site}(*)} \rightarrow \Omega^{\cdot}(*)[[z]]$$

is a quasi-isomorphism by Lemma 3.2 □

3.2.8 It follows from Lemma 3.8 by applying $p^* \circ Ri$ that

$$p^* \circ Ri \circ i^\sharp \circ Rg_* \circ Ri(i^\sharp \mathbb{R}_{\mathbf{Site}([*/S^1])}) \cong p^* \circ Ri \circ i^\sharp \mathbb{R}[[z]]_{\mathbf{Site}(*)}.$$

We now use the second assertion of Lemma 2.50 in order to commute $Ri \circ i^\sharp$ with p^* . We get

$$Ri \circ i^\sharp \circ p^* \circ Rg_* \circ Ri(i^\sharp \mathbb{R}_{\mathbf{Site}([*/S^1])}) \cong Ri \circ i^\sharp \circ p^* \mathbb{R}[[z]]_{\mathbf{Site}(*)}.$$

We now apply i^\sharp and use that $i^\sharp \circ Ri \cong \text{id}$ in order to drop the functor Ri and get the quasi-isomorphism

$$i^\sharp \circ p^* \circ Rg_* \circ Ri(i^\sharp \mathbb{R}_{\mathbf{Site}([*/S^1])}) \cong i^\sharp \circ p^* \mathbb{R}[[z]]_{\mathbf{Site}(*)}.$$

By the explicite description of p^* given in the proof of Lemma 2.9 we see that

$$p^*(\mathbb{R}[[z]]_{\text{Site}(\ast)}) \cong \mathbb{R}[[z]]_{\mathbf{X}} .$$

We thus have a quasi-isomorphism

$$i^\# \circ p^* \circ Rg_* \circ Ri(i^\# \mathbb{R}_{\text{Site}(\ast/S^1)}) \cong i^\#(\mathbb{R}[[z]]_{\mathbf{X}}) . \quad (3.9)$$

Combining the isomorphisms (3.6), (3.7) and (3.9) we obtain a quasi-isomorphism

$$i^\# C_A(\Omega(G)) \cong i^\#(\mathbb{R}[[z]]_{\mathbf{X}}) .$$

In particular we see that z generates the cohomology.

3.2.9 Since X is contractible we find $\gamma \in \Omega^2(X)$ such that $d_{dR}\gamma = \lambda$. We define a map of complexes of sheaves

$$\psi : i^\# \mathbb{R}[[z]]_{\mathbf{X}} \rightarrow \Omega[[z]]_0 \xrightarrow{e^{-\gamma T}} \Omega[[z]]_\lambda .$$

The first map is given by the inclusion $i^\# \mathbb{R}_{\mathbf{X}} \rightarrow \Omega_X$ and is a quasi-isomorphism. The second map is an isomorphism of sheaves of complexes. Therefore ψ is a quasi-isomorphism. Note that ψ is multiplicative and $\psi(z) = z - \gamma$. We further define $\kappa : i^\# \mathbb{R}[[z]]_{\mathbf{X}} \rightarrow i^\# C_A(\Omega(G))$ such that $\kappa(z) = (\alpha, \beta - \gamma) = \phi(z - \gamma)$. Then we have a commutative diagram

$$\begin{array}{ccc} i^\# \mathbb{R}[[z]]_{\mathbf{X}} & \xrightarrow{\kappa} & i^\# C_A(\Omega(G)) \\ & \searrow \psi & \nearrow \phi \\ & \Omega[[z]]_\lambda & \end{array} .$$

If we show that κ is a quasi-isomorphism, then since ψ is a quasi-isomorphism, ϕ must be a quasi-isomorphism, too. It suffices to see that $\kappa(z) := (\alpha, \beta - \gamma)$ represents a non-trivial cohomology class. Assume that it is a boundary locally on $(U \rightarrow X) \in \mathbf{X}$. Then there exists $x \in \Omega^0(G)(A_U^1)$ and $y \in \Omega^1(G)(A_U^0)$ such that $\delta x = 0$, $d_{dR}x + \delta y = \alpha_U$ and $d_{dR}y = (\beta - \gamma)_U$ (the subscript indicates that the forms are pulled back to A_U^*). By exactness of the δ -complex we can in fact assume that $x = 0$. But then the equation $\delta y = \alpha_U$ is impossible since δy vanishes on vertical vectors on the bundles $A_U^1 \rightarrow A_U^0$ given by the source and range projections while α as a connection form is non-trivial on those vectors. \square

3.2.10 We now finish the proof of Theorem 1.1. We combine Proposition 3.4 with 3.1.4 and the fact that f_* preserves sheaves (Lemma 2.13) in order to get

$$i^\# \circ Rf_* \circ Ri(i^\# \mathbb{R}_{\mathbf{G}}) \cong i^\# C_A(\Omega(G)) \cong \Omega[[z]]_\lambda .$$

\square

References

- [1] Michael Atiyah and Graeme Segal. Twisted K-theory and cohomology. 1.3.1, 1.3.3, 1.3.5, (2)
- [2] Michael Atiyah and Graeme Segal. Twisted K -theory. *Ukr. Mat. Visn.*, 1(3):287–330, 2004. 1.3.6
- [3] Peter Bouwknegt, Alan L. Carey, Varghese Mathai, Michael K. Murray, and Danny Stevenson. Twisted K -theory and K -theory of bundle gerbes. *Comm. Math. Phys.*, 228(1):17–45, 2002. 1.3.1, 1.3.5

- [4] Peter Bouwknegt, Jarah Evslin, and Varghese Mathai. *T*-duality: topology change from *H*-flux. *Comm. Math. Phys.*, 249(2):383–415, 2004. 1.3.1
- [5] Jean-Luc Brylinski. *Loop spaces, characteristic classes and geometric quantization*, volume 107 of *Progress in Mathematics*. Birkhäuser Boston Inc., Boston, MA, 1993. 1.3.9, 1.3.14, (3)
- [6] U. Bunke, Th. Schick, and M. Spitzweck. Foundations of sheaf theory on topological stacks. In preparation. 1.1.3, 2.2.5
- [7] U. Bunke, Th. Schick, and M. Spitzweck. Inertia and delocalized twisted cohomology. In preparation. 1.1.3
- [8] U. Bunke, Th. Schick, and M. Spitzweck. *T*-duality and periodic twisted cohomology. In preparation. 1.1.3
- [9] Daniel S. Freed, Michael J. Hopkins, and Constantin Teleman. Twisted equivariant K-theory with complex coefficients. 1.3.3
- [10] Jochen Heinloth. Survey on topological and smooth stacks. In *Mathematisches Institut Göttingen, WS04-05 (Y. Tschinkel, ed.)*, pages 1–31. 2005. 1.3.9, 1.3.12, 1.3.14, 2.2.1, (4), (5), 2.2.5
- [11] Nigel Hitchin. Lectures on special Lagrangian submanifolds. In *Winter School on Mirror Symmetry, Vector Bundles and Lagrangian Submanifolds (Cambridge, MA, 1999)*, volume 23 of *AMS/IP Stud. Adv. Math.*, pages 151–182. Amer. Math. Soc., Providence, RI, 2001. 1.3.7, 1.3.8, 3.2.1, 3.2.2
- [12] Michael Joachim. Higher coherences for equivariant *K*-theory. In *Structured ring spectra*, volume 315 of *London Math. Soc. Lecture Note Ser.*, pages 87–114. Cambridge Univ. Press, Cambridge, 2004. 1.3.6
- [13] Masaki Kashiwara and Pierre Schapira. *Sheaves on manifolds*, volume 292 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, 1994. With a chapter in French by Christian Houzel, Corrected reprint of the 1990 original. 3.1.2
- [14] Gérard Laumon and Laurent Moret-Bailly. *Champs algébriques*, volume 39 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics*. Springer-Verlag, Berlin, 2000. 1.3.19, 2.2.11
- [15] Varghese Mathai and Danny Stevenson. On a generalized Connes-Hochschild-Kostant-Rosenberg theorem. *Advances in Mathematics*, vol. 200 no. 2 (2006) 1-33. 1.3.1, 1.3.5
- [16] Varghese Mathai and Danny Stevenson. Chern character in twisted *K*-theory: equivariant and holomorphic cases. *Comm. Math. Phys.*, 236(1):161–186, 2003. 1.3.1, 1.3.5
- [17] J. P. May and J. Sigurdsson. Parametrized homotopy theory. 1.3.4
- [18] J. Peter May. *E_∞ ring spaces and E_∞ ring spectra*. Springer-Verlag, Berlin, 1977. With contributions by Frank Quinn, Nigel Ray, and Jørgen Tornehave, Lecture Notes in Mathematics, Vol. 577. 1.3.5
- [19] David Metzler. Topological and Smooth Stacks. 2.2.1, 2
- [20] M. K. Murray. Bundle gerbes. *J. London Math. Soc. (2)*, 54(2):403–416, 1996. 1.3.7
- [21] Michael K. Murray and Daniel Stevenson. Bundle gerbes: stable isomorphism and local theory. *J. London Math. Soc. (2)*, 62(3):925–937, 2000. 1.3.7
- [22] Behrang Noohi. Foundations of Topological Stacks I. 1.3.9, 1.3.14, 2.2.1
- [23] Martin Olsson. Sheaves on artin stacks. 2.2.11

- [24] Günter Tamme. *Introduction to étale cohomology*. Universitext. Springer-Verlag, Berlin, 1994. Translated from the German by Manfred Kolster. 2.1.1, 2.1.3, 2.2.6, 2.2.9, 2.2.10, 2.2.10, 2.2.11, 2.3.2, 2.3.4, 2.3.4, 2.28, 2.3.7, 2.4.7, 2.5.6, 3.1.2
- [25] Jean-Louis Tu, Ping Xu, and Camille Laurent-Gengoux. Twisted k-theory of differentiable stacks. 1.3.8