

FINITENESS OF SPATIAL CENTRAL CONFIGURATIONS IN THE FIVE-BODY PROBLEM

MARSHALL HAMPTON AND ANDERS JENSEN

ABSTRACT. We strengthen a generic finiteness result due to Moeckel by showing that the number of spatial central configurations of the Newtonian five-body problem with positive masses is finite, apart from some explicitly given special cases of mass values.

1. INTRODUCTION

In this paper we present a computer-assisted proof of the finiteness of the spatial central configurations of the Newtonian five-body problem with positive masses, with the exception of some explicit special cases of mass values.

By the Newtonian spatial n -body problem we mean the dynamical system given by

$$(1) \quad m_j \ddot{x}_j = \sum_{i \neq j} \frac{m_i m_j (x_i - x_j)}{r_{ij}^3} \quad 1 \leq j \leq n$$

where $x_i \in \mathbb{R}^3$ is the position of particle i , r_{ij} is the distance between x_i and x_j , and m_i is the mass of particle i [14].

A central configuration of the n -body satisfies the equations

$$(2) \quad \lambda(x_j - c) = \sum_{i \neq j} \frac{m_i (x_i - x_j)}{r_{ij}^3} \quad 1 \leq j \leq n$$

where $\lambda < 0$ and c is the center of mass. Such a configuration, if started from rest, will collapse to the center of mass. In the planar case these configurations also give rise to relative equilibria - orbits in which each particle moves on a circle at a common angular speed.

For the Newtonian three-body and four-body problems it is known that central configurations in dimensions one, two, and three are finite for positive masses [4, 5, 9, 13]. There is also a generic finiteness result

1991 *Mathematics Subject Classification.* 70F10, 70F15, 37N05, 76Bxx.

Key words and phrases. Celestial mechanics, n -body problem, tropical geometry.

for ‘Dziobek configurations’ which applies to the case we are studying [11] since the dimension of our configurations is $n - 2$.

The five-body problem is of particularly interest in this context because of the known continuum of planar central configurations for the five-body problem if one negative mass is permitted [15]. Recently Alain Albouy and Vadim Kaloshin have proved that the planar five-body central configurations are finite apart from some explicitly given special cases [2].

We have proven (with computer assistance on exact computations) the following result:

Theorem 1.1. *There are finitely many central configurations of the Newtonian spatial five-body problem with the possible exception of some mass parameter values which vanish on an explicitly given set of polynomials.*

The proof of this theorem is given by the remainder of our paper. In Section 2 we briefly review the necessary tropical geometry. Following that we describe the equations we used in Section 3. The exceptional cases are given in Table 1 of Section 4 along with a description of the tropical prevariety of our equations. The next two sections (5 and 6) give details of the two most complicated cases that arose from the tropical prevariety.

2. TROPICAL GEOMETRY

Our proof strategy will be the same as that of [5], but we will use the language of tropical geometry. Our equations for central configurations (described in Section 3) define an algebraic variety in the algebraic torus $(\mathbb{C}^*)^{10}$. Instead of proving that the algebraic variety has dimension 0 we will attempt to prove that its tropical variety is a polyhedral fan of dimension 0. Both varieties depend on the choice of masses. We will consider these masses as being unknown positive real numbers. In this section we give the necessary definitions in tropical geometry and refer to [20, Chapter 9] for a general introduction.

Definition 2.1. *Let $I \subseteq \mathbb{C}[x_1, \dots, x_n]$ be an ideal and $\omega \in \mathbb{R}^n$ a vector. For a monomial $x^v := x_1^{v_1} \cdots x_n^{v_n}$ with $v \in \mathbb{N}^n$ we define its ω -degree as $\omega \cdot v$. The initial form $\text{in}_\omega(f)$ of a polynomial $f \in \mathbb{C}[x_1, \dots, x_n]$ is the sum of all terms of f with maximal ω -degree. The initial ideal of I with respect to ω is defined as $\text{in}_\omega(I) := \langle \text{in}_\omega(f) : f \in I \rangle$.*

We note that the initial ideal is not always a monomial ideal.

Definition 2.2. *The tropical variety of an ideal $I \subseteq \mathbb{C}[x_1, \dots, x_n]$ is the set*

$$T(I) := \{\omega \in \mathbb{R}^n : \text{in}_\omega(I) \text{ does not contain a monomial}\}.$$

For an ideal $J \subseteq \mathbb{C}[x_1, \dots, x_n]$ and a polynomial $f \in \mathbb{C}[x_1, \dots, x_n]$ we define the *saturation* $(J : f^\infty) := \{g \in \mathbb{C}[x_1, \dots, x_n] : \exists m : gf^m \in J\}$. The saturation $(\text{in}_\omega(I) : x_1 \cdots x_n^\infty)$ is $\langle 1 \rangle$ if and only if $\omega \notin T(I)$.

The tropical variety $T(I)$ can be thought of as a polyhedral fan. In the case where I is homogeneous, $T(I)$ is the support of a subfan of the *Gröbner fan* of I . The Gröbner fan of I consists of all cones which are closures of maximal sets of vectors in \mathbb{R}^n giving rise to the same initial ideal of I .

The following theorem was first stated by Bieri and Groves in terms of valuations. A proof in the language of initial ideals was given in [20], see also [7, Chapter 8]. It will be essential for our arguments.

Theorem 2.3 (Bieri Groves). *Let $I \subseteq \mathbb{C}[x_1, \dots, x_n]$ be a prime ideal and let d be the dimension of the variety $V(I) \subseteq (\mathbb{C}^*)^n$ defined by I . The tropical variety $T(I)$ is the support of a pure d -dimensional polyhedral fan.*

It is not difficult to prove that if $I = \bigcap_i Q_i$ then $T(I) = \bigcup T(\sqrt{Q_i})$, see for example [7]. Therefore, if I is not a prime ideal, we can apply the theorem to a primary decomposition of I to get the largest dimension of a cone in $T(I)$.

For our purpose it will suffice to compute $T(I)$ and see that it is zero-dimensional to conclude that $V(I)$ is zero-dimensional. A first approximation to $T(I)$ comes from the following description, see [20].

Proposition 2.4. *Let $I \subseteq \mathbb{C}[x_1, \dots, x_n]$. The following identity holds*

$$T(I) = \bigcap_{f \in I} T(\langle f \rangle).$$

Given a set of generators f_1, \dots, f_m of I we may compute the *tropical prevariety* $\bigcap_i T(\langle f_i \rangle)$ defined by f_1, \dots, f_m . However, the prevariety does not equal $T(I)$ in general. The software Gfan [8] will compute the prevariety as a polyhedral fan. For each cone in this fan of positive dimension we wish to check if the cone is contained in $T(I)$. It is possible that some relative interior points are in $T(I)$ and others are not. To check if a single ray ω is contained in $T(I)$ we can in theory compute $\text{in}_\omega(I)$. However, computing initial ideals is usually done by computing a Gröbner basis ([19, Corollary 1.9]). This computation is not feasible in our setting. Indeed, if we could compute a Gröbner basis for I we would also know the dimension of I .

Instead we will approximate $\text{in}_\omega(I)$ by choosing a big generating set for I and taking initial forms with respect to ω . If the ideal generated by these initial forms contains a monomial, then so does $\text{in}_\omega(I)$.

We need two more properties of tropical varieties. First, tropical varieties are balanced which implies that a tropical variety in 1-dimensional space \mathbb{R} is either empty, the origin or all of \mathbb{R} . Second, a rational projection of a tropical variety to a linear subspace is again a tropical variety. This follows from [6, Theorem 3.1] and the possibility to do multiplicative changes of coordinates in the Laurent polynomial ring. Putting this together we get that to show that a tropical variety $T(I)$ equals the origin, it suffices to show that $T(I) \cap \{\omega \in \mathbb{R}^n : \sum_i \omega_i \geq 0\} = \{0\}$.

3. EQUATIONS FOR THE SPATIAL CASE

We will use two versions of the Albouy-Chenciner equations for central configurations (originally described in [1]). The first version is identical to that discussed in [5], namely the polynomial versions (with denominators cleared) of

$$(3) \quad f_{ij} = \sum_{k=1}^n m_k [S_{ik}(r_{jk}^2 - r_{ik}^2 - r_{ij}^2) + S_{jk}(r_{ik}^2 - r_{jk}^2 - r_{ij}^2)] = 0$$

where $S_{ij} = r_{ij}^{-3} + \lambda$. We choose to set $\lambda = -1$, which can be done without loss of generality because of the homogeneity of the equations above. We will refer to these as the symmetric Albouy-Chenciner equations. We will denote the complete set of the f_{ij} as \mathcal{F} .

Gareth Roberts observed [16] that a more restrictive set of equations can be obtained from the Albouy-Chenciner linear operator equations, namely

$$(4) \quad g_{ij} = \sum_{k=1}^n m_k S_{ik}(r_{jk}^2 - r_{ik}^2 - r_{ij}^2) = 0.$$

Since $g_{ij} \neq g_{ji}$, these give 20 equations in the five-body problem. Since $f_{ij} = g_{ij} + g_{ji}$ the f_{ij} equations are redundant but they were included in the tropical intersection calculation anyway in order to generate a more refined set of cones. These equations will collectively be denoted as \mathcal{G} .

Fifteen more ‘Dziobek’ equations were added [3]. Only five of these are independent, but all were included to preserve symmetry:

$$(5) \quad h_{ijkl} = (r_{ij}^{-3} - 1)(r_{kl}^{-3} - 1) - (r_{ik}^{-3} - 1)(r_{jl}^{-3} - 1) = 0$$

where i, j, k , and l are distinct indices. The set of Dziobek equations will be denoted \mathcal{H} .

The distances between five points in \mathbb{R}^3 satisfy a single constraint, that the determinant of the Cayley-Menger matrix is zero:

$$(6) \quad e_{CM} = \det \begin{pmatrix} 0 & 1 & 1 & 1 & 1 & 1 \\ 1 & 0 & r_{12}^2 & r_{13}^2 & r_{14}^2 & r_{15}^2 \\ 1 & r_{12}^2 & 0 & r_{23}^2 & r_{24}^2 & r_{25}^2 \\ 1 & r_{13}^2 & r_{23}^2 & 0 & r_{34}^2 & r_{35}^2 \\ 1 & r_{14}^2 & r_{24}^2 & r_{34}^2 & 0 & r_{45}^2 \\ 1 & r_{15}^2 & r_{25}^2 & r_{35}^2 & r_{45}^2 & 0 \end{pmatrix} = 0$$

which we included in our set of equations.

Central configurations satisfy the equation $U/I = -\lambda M$, where I is the moment of inertia $I = \frac{1}{M} \sum_{i < j} m_i m_j r_{ij}^2$, $M = \sum m_i$, $U = \sum_{i < j} m_i m_j r_{ij}^{-1}$ (see for example [12]). Since we chose $\lambda = -1$ we have the equation

$$(7) \quad e_{IU} = U - MI = 0$$

This is redundant, as it is a consequence of the Albouy-Chenciner equations, but was included (in polynomial form, with cleared denominators) since we thought its simplicity and symmetry might be helpful in the analysis of the tropical prevariety.

In summary, in what follows we will be analyzing the ideal $I_s := \langle \mathcal{F}, \mathcal{G}, \mathcal{H}, e_{CM}, e_{IU} \rangle$.

4. THE TROPICAL PREVARIETY

We calculated the tropical prevariety of the set of 42 equations described in the previous section, using Gfan [8]. There were 576 rays, with 26 distinct types after considering the symmetry of the equations. In addition, there are 50 distinct cases of two-dimensional cones, 27 distinct cases of three-dimensional cones, 11 distinct cases of four-dimensional cones, and 3 distinct cases of five-dimensional cones.

Most of these cases can be easily dismissed by computing a Gröbner basis of the initial forms and saturating with respect to the r_{ij} and m_i variables. In many cases, if we eliminate the r_{ij} after saturating we obtain a sum over some subset of the masses (e.g. $m_1 + m_2 + m_3$) within the elimination ideal, which means there are no positive mass solutions within that cone. However, for a few rays and one-dimensional cones there were some nonzero solutions for some choice of masses. Gröbner bases were calculated using Singular [18] and Sage [17].

Representatives of these exceptional cases are given in the following table along with the indices of the rays generating the cone and their directions. All polynomials listed for a given cone must be satisfied.

Ray indices	Interior point	Exceptional polynomials
[59]	(1, 0, 0, 0, 0, 0, 1, 1, 0)	$m_1m_2 - m_3m_4 - m_3m_5$
[72]	(1, 0, 0, 0, 0, 0, 1, 1, 1)	$(4m_1m_2 - (m_3 + m_4)^2 - (m_3 + m_5)^2 + m_3^2)$
[59, 72]	(2, 0, 0, 0, 0, 0, 2, 2, 1)	$m_3 - m_4 - m_5,$ $m_4^2 + 2m_4m_5 + m_5^2 - m_2m_1$
[193, 210]	(2, 2, -2, -2, 0, 2, 0, 2, 0, -2)	$m_1 - m_4, m_2 - m_3$
[210, 453]	(4, 4, 0, 0, 0, 4, 2, 4, 2, 0)	$m_1 - m_4, m_2 - m_3$
[268, 453]	(4, 4, 0, 0, 0, 4, 3, 4, 3, 0)	$m_1 - m_4, m_2 - m_3$
[270]	(1, 1, 1, 0, 0, 0, 1, 0, 1, 1)	see section 5
[275]	(1, 1, 1, 0, 1, 1, 0, 1, 0, 0)	see section 6

TABLE 1. Representatives of exceptional cases

5. A TROUBLESOME RAY

In this section we explain the possible exceptions to the finiteness result stemming from the ray with weight $\omega_{270} = (1, 1, 1, 0, 0, 1, 0, 1, 1)$.

Recall that $I_s = \langle \mathcal{F}, \mathcal{G}, \mathcal{H}, e_{CM}, e_{IU} \rangle$. This is an ideal in $\mathbb{R}[r_{12}, \dots, r_{45}]$. Our goal is to show that $J := (\text{in}_{\omega_{270}}(I_s) : r_{12} \cdots r_{45}^\infty)$ contains a monomial, or equivalently, that it is equal to $\langle 1 \rangle$.

We first consider the initial forms of the 15 Dziobek equations, \mathcal{H} . Six of these are binomials. Among these are $r_{12}^3 r_{13}^3 r_{34}^3 - r_{12}^3 r_{13}^3 r_{24}^3$ and $r_{13}^3 r_{14}^3 r_{24}^3 - r_{13}^3 r_{14}^3 r_{23}^3$. We conclude that $r_{34}^3 - r_{24}^3$ and $r_{24}^3 - r_{23}^3$ are in J . Similarly, the initial form $-r_{13}^3 r_{25}^3 + r_{13}^3 r_{23}^3 r_{25}^3 + r_{13}^3 r_{15}^3 r_{25}^3$ shows that $r_{23}^3 + r_{15}^3 - 1$ is in J .

We now define $J' = \langle \text{initial forms of } (\mathcal{F}, \mathcal{G}, e_{CM}, e_{IU}) + \langle r_{34}^3 - r_{24}^3, r_{24}^3 - r_{23}^3, r_{23}^3 + r_{15}^3 - 1 \rangle \subseteq J$. Performing the substitutions $r_{23}^3 \mapsto x, r_{24}^3 \mapsto x, r_{34}^3 \mapsto x, r_{15}^3 \mapsto 1 - x$ on the generators of J' and removing monomial factors, we get a new ideal whose generators do not contain r_{23}, r_{24}, r_{34} and r_{15} . Thus instead of considering J' , we consider $\tilde{J}' \subseteq \mathbb{R}[r_{13}, r_{14}, r_{25}, r_{35}, r_{45}, x]$. This ideal is easy to handle. We compute

$$\begin{aligned} & (\tilde{J}' : r_{12}r_{13}r_{14}r_{25}r_{35}r_{45}x(m_1 + m_5)^\infty) \cap \mathbb{Q}[m_1, m_5, r_{13}, r_{14}, r_{35}, r_{45}] \\ &= \langle m_1r_{14}^2 + m_5r_{45}^2, m_1r_{13}^2 + m_5r_{35}^2, r_{14}^2r_{35}^2 - r_{13}^2r_{45}^2 \rangle, \end{aligned}$$

where we regard \tilde{J}' as an ideal in $\mathbb{Q}[m_1, \dots, m_5, r_{13}, r_{14}, r_{25}, r_{35}, r_{45}, x]$. For every positive real choice of masses, we conclude that $m_1r_{14}^2 +$

$m_5 r_{45}^2, m_1 r_{13}^2 + m_5 r_{35}^2, r_{14}^2 r_{35}^2 - r_{13}^2 r_{45}^2 \in J$. Similarly, by choosing different subrings for the intersection we get that $\{m_1 r_{12}^2 + m_5 r_{25}^2, r_{14}^2 r_{25}^2 - r_{12}^2 r_{45}^2, r_{13}^2 r_{25}^2 - r_{12}^2 r_{35}^2\} \subseteq J$.

Now, define $K = \langle m_1 r_{12}^2 + m_5 r_{25}^2, m_1 r_{14}^2 + m_5 r_{45}^2, m_1 r_{13}^2 + m_5 r_{35}^2, r_{14}^2 r_{35}^2 - r_{13}^2 r_{45}^2, r_{14}^2 r_{25}^2 - r_{12}^2 r_{45}^2, r_{13}^2 r_{25}^2 - r_{12}^2 r_{35}^2 \rangle$. Let d_1, d_2, d_3 be the initial forms of the Dziobek equations which are not binomial or trinomials – there are only three up to sign. The ideals $(\langle d_i \rangle \cap \mathbb{Q}[m_1, m_5, r_{12}, r_{13}, r_{14}]) : r_{12} r_{13} r_{14} (m_1^3 + m_5^3)^\infty$ equal

$$\begin{aligned} & \langle (r_{12}^3 - r_{14}^3)(m_1(r_{12}^3 - r_{14}^3)^2 + m_5(r_{12}^3 + r_{14}^3)^2) \rangle \\ & \langle (r_{13}^3 - r_{14}^3)(m_1(r_{13}^3 - r_{14}^3)^2 + m_5(r_{13}^3 + r_{14}^3)^2) \rangle \\ & \langle (r_{12}^3 - r_{13}^3)(m_1(r_{12}^3 - r_{13}^3)^2 + m_5(r_{12}^3 + r_{13}^3)^2) \rangle. \end{aligned}$$

We conclude that each of these generators, denoted k_i , are in J for every choice of positive masses. Similarly, $(\langle \tilde{J}' \cap \mathbb{Q}[r_{12}, r_{13}, r_{14}, r_{12}] \rangle : (m_1 + m_5)^\infty) = \langle m_2 r_{12}^2 + m_3 r_{13}^2 + m_4 r_{14}^2 \rangle$ shows that J contains $m_2 r_{12}^2 + m_3 r_{13}^2 + m_4 r_{14}^2$ for every positive choice of masses.

So it suffices to consider the ideal $\langle m_2 r_{12}^2 + m_3 r_{13}^2 + m_4 r_{14}^2, k_1, k_2, k_3 \rangle$ homogeneous in $r_{12}, r_{13},$ and r_{14} , homogeneous in m_1 and m_5 , and homogeneous in $m_2, m_3,$ and m_4 . If we dehomogenize by setting $r_{14} = 1$ and $m_5 = 1$ we obtain the following:

$$(8) \quad (r_{12}^3 - 1)(r_{12}^6 + 2\frac{1 - m_1^3}{1 + m_1^3}r_{12}^3 + 1) = 0$$

$$(9) \quad (r_{13}^3 - 1)(r_{13}^6 + 2\frac{1 - m_1^3}{1 + m_1^3}r_{13}^3 + 1) = 0$$

$$(10) \quad (r_{12}^3 - r_{13}^3)(r_{12}^6 + 2\frac{1 - m_1^3}{1 + m_1^3}r_{12}^3 r_{13}^3 + r_{13}^6) = 0$$

$$(11) \quad r_{12}^2 m_2 + r_{13}^2 m_3 + m_4 = 0$$

Equations (8) and (9) imply that both r_{12} and r_{13} lie on the unit circle at one of nine points, either at a root of unity or at one of the six locations given by the formula below, in which the choice of cube root is unspecified:

$$\left(\frac{m_1^3 - 1 \pm 2I\sqrt{m_1^3}}{m_1^3 + 1} \right)^{1/3}.$$

These roots are all distinct for positive m_1 .




FIGURE 1. Exceptional cases for ray 270 in the plane $m_2 + m_3 + m_4 = 1$.

Equation (10) imposes the further constraint that the ratio of the roots chosen for r_{12} and r_{13} must also be equal to one of the nine points described above; this in turn means that at least one of r_{12} and r_{13} must be a complex root of unity. In addition, the positivity of the masses combined with equation (11) means that the imaginary parts of r_{12}^2 and r_{13}^2 must have opposite signs.

The consequence of this is that for each choice of m_1/m_5 there are finitely many choices for m_2/m_4 and m_3/m_4 . By choosing both r_{12} and r_{13} to be roots of unity with opposing imaginary parts, we always have $m_2 = m_3 = m_4$ as a solution. In Figure 1 we show the other possibilities by normalizing $m_2 + m_3 + m_4 = 1$ and projecting the set of positive mass solutions orthogonally onto the plane:

It is straightforward to obtain an explicit polynomial in the masses by eliminating the distance variables, but this polynomial is rather large and unwieldy; we believe the structure of the real solutions is much clearer in the above system.

6. AN EVEN MORE TROUBLESOME RAY

In this section we explain the possible exceptions to the finiteness result stemming from the ray with weight $\omega_{275} = (1, 1, 1, 0, 1, 1, 0, 1, 0, 0)$.

The initial forms of three of the Dziobek equations give us equations of the form

$$r_{ij}^{-3} + r_{kl}^{-3} - r_{ik}^{-3} - r_{jl}^{-3} = 0$$

for all choices $\{i, j, k, l\} = \{1, 2, 3, 4\}$.

After removing common monomial factors, the remaining equations are binomials of the form $r_{r_5}^3 - r_{s_5}^3$. Using these binomials we may rewrite the Albouy-Chenciner initial form $\text{in}_{\omega_{275}}(f_{i5})$ as $\sum_{j \notin \{i, 5\}} m_j r_{ij}^2$. This works for $i = 1, \dots, 4$.

Taking these four equations and multiplying the Dziobek equations by monomials, we get the following ideal in $\mathbb{Q}[m_1, \dots, m_4, r_{13}, \dots, r_{34}]$:

$$\begin{aligned} J' := & \langle m_4 r_{14}^2 + m_3 r_{13}^2 + m_2 r_{12}^2, m_4 r_{24}^2 + m_3 r_{23}^2 + m_1 r_{12}^2, \\ & m_4 r_{34}^2 + m_2 r_{23}^2 + m_1 r_{13}^2, m_3 r_{34}^2 + m_2 r_{24}^2 + m_1 r_{14}^2, \\ & -r_{13}^3 r_{24}^3 r_{34}^3 + r_{12}^3 r_{24}^3 r_{34}^3 + r_{12}^3 r_{13}^3 r_{34}^3 - r_{12}^3 r_{13}^3 r_{24}^3, \\ & -r_{14}^3 r_{23}^3 r_{24}^3 + r_{13}^3 r_{23}^3 r_{24}^3 + r_{13}^3 r_{14}^3 r_{24}^3 - r_{13}^3 r_{14}^3 r_{23}^3, \\ & -r_{14}^3 r_{23}^3 r_{34}^3 + r_{12}^3 r_{23}^3 r_{34}^3 + r_{12}^3 r_{14}^3 r_{34}^3 - r_{12}^3 r_{14}^3 r_{23}^3 \rangle \end{aligned}$$

The initial form of the Cayley-Menger determinant e_{CM} has four factors:

$$\begin{aligned} & (-r_{14}r_{23} - r_{13}r_{24} + r_{12}r_{34})(-r_{14}r_{23} + r_{13}r_{24} + r_{12}r_{34}) \\ & (r_{14}r_{23} - r_{13}r_{24} + r_{12}r_{34})(r_{14}r_{23} + r_{13}r_{24} + r_{12}r_{34}) = 0 \end{aligned}$$

The first three of these factors are equivalent under permutations of $(1, 2, 3, 4)$, so we will only analyze the cases in which the first or last factor vanish.

We consider the ideals $J' + \langle r_{14}r_{23} + r_{13}r_{24} + r_{12}r_{34} \rangle$ and $J' + \langle r_{14}r_{23} + r_{13}r_{24} - r_{12}r_{34} \rangle$. We wish to saturate with respect to all variables and thereafter eliminate the r -variables. Since our ideals are homogeneous in the grading of the masses and in the grading of the r -variables, any reduced Gröbner basis will work as an elimination Gröbner basis and our difficulty will be to saturate the variables rather than eliminating. The saturation can be carried out in Singular using for example [19, Lemma 12.1] for computing saturations by reverse lexicographic term orders. Saturating with respect to the variables in the following order worked out for us:

$$m_4, m_3, r_{34}, r_{24}, r_{23}, r_{14}, r_{12}, r_{13}, m_1, m_2.$$

In each case after computing the elimination ideal, we are left with a single polynomial in the cubes of the masses. For simplicity we will

use $M_i = m_i^3$ as variables. The polynomial is homogeneous of degree 12, with a maximum degree of 6 in each variable. For the case in which $-r_{14}r_{23} - r_{13}r_{24} + r_{12}r_{34} = 0$ we will denote this polynomial by P_1 , and in the case $r_{14}r_{23} + r_{13}r_{24} + r_{12}r_{34}$ the polynomial will be denoted by P_2 .

Lemma 6.1. *The only real positive solutions for P_1 are when $M_1 = M_2$ (which implies $m_1 = m_2$ for real masses) and*

$$\begin{aligned} P_{34} = & M_1^4 - 2M_1^3M_3 - 2M_1^3M_4 + M_1^2M_3^2 - 12M_1^2M_3M_4 + M_1^2M_4^2 \\ & - 2M_1M_3^2M_4 - 2M_1M_3M_4^2 + M_3^2M_4^2 = 0 \end{aligned}$$

or the corresponding case with $\{M_1, M_2\}$ interchanged with $\{M_3, M_4\}$.

Proof. We will prove the result using interval arithmetic. Because P_1 is homogeneous and symmetric under the interchanges $M_1 \leftrightarrow M_2$, $M_3 \leftrightarrow M_4$, and $(M_1, M_2) \leftrightarrow (M_3, M_4)$, we can dehomogenize by letting $M_4 = 1 - M_1 - M_2 - M_3$ and restrict to the set Ω given by $0 \leq M_1 \leq M_2 \leq 1 - M_1 - M_2 - M_3$ and $M_3 \leq 1 - M_1 - M_2 - M_3$. It was convenient to decompose this set into three tetrahedra Ω_A , Ω_B , and Ω_C with $M_2 \leq M_3$, $M_1 \leq M_3 \leq M_2$, and $M_3 < M_1$ respectively.

Our goal is to prove that $P_1 > 0$ in the interior of Ω . Our strategy is to show that the only critical points of P_1 in the interior of Ω are positive. P_1 factors nicely on the boundary of Ω and there it is easy to see that there are the zero-valued curves of minima described in the statement of the lemma.

To examine the critical points in Ω with interval arithmetic is somewhat challenging due to the degeneracy of the zero set on its boundary, so we first applied the linear transformation A taking M -coordinates to y -coordinates according to the identities $M_1 = y_1/4$, $M_2 = y_2/3 + y_1/4$, $M_3 = 1/2 - y_1/4 - y_2/6 - y_3/2$. The tetrahedron Ω_A maps to the tetrahedron $\hat{\Omega}_A$ given by $y_i \geq 0$, $y_1 + y_2 + y_3 \leq 1$. The key advantage of this transformation is that the zero set of $\hat{P}_1(y_1, y_2, y_3) = P_1(A^{-1}(y_1, y_2, y_3))$ is contained in the planes $y_2 = 0$ and $y_3 = 0$. This property causes the resultants $R_{12} = \text{Res}_{y_1}(\partial \hat{P}_1 / \partial y_1, \partial \hat{P}_1 / \partial y_2)$ and $R_{23} = \text{Res}_{y_1}(\partial \hat{P}_1 / \partial y_2, \partial \hat{P}_1 / \partial y_3)$ in $\mathbb{C}[y_2, y_3]$ to have monomial factors of high degree instead of more complicated polynomial factors.

Some of the factors of these resultants can be seen to have no zeros in the projection of $\hat{\Omega}$ to the (y_2, y_3) plane. For example, R_{12} factorizes as

$$y_2^5 y_3^4 (y_2 - 3y_3 - 3)^2 (y_2 - 3y_3 + 3)^2 (y_2 + 3y_3 - 3)^2 (y_2 + 3y_3 + 3)^2 (y_2^2 + 9y_3^2 - 9)^2 f_1 f_2$$

where f_1 has degree 24 and f_2 has degree 48. It is straightforward to show that all the factors other than f_1 and f_2 do have no zeros in $\hat{\Omega}$.

We use recursive interval arithmetic applied to \hat{P}_1 , the derivatives of \hat{P}_1 , and factors of the above resultants to exclude the existence of non-positive critical values in the interior of Ω except for very small neighborhoods of the two critical points $(M_1, M_2, M_3) \in \{(0, 0, 0), (1/3, 1/3, 0)\}$ where P_1 is zero. For a given polynomial in n -variables x_1, \dots, x_n we first view it in the ring $\mathbb{Q}[x_1, \dots, x_{n-1}][x_n]$ and obtain lower and upper bounds to the coefficients (which are single-variable polynomials in x_n) by verifying numerical estimates with the real root isolation code of Carl Witty in Sage [21]. The remaining very small neighborhoods can be excluded by considering expansions of the second derivatives of P_1 around these points. \square

P_2 is simpler than P_1 since it is symmetric under any interchange of masses. Thus it suffices to examine P_2 on the set Ω_A . The only real positive solutions occur when three masses are equal.

It should be possible to obtain much simpler proofs of the above statements by finding an appropriate decomposition of P_1 and P_2 , but we were unable to do so.

7. CONCLUSION

We have shown that the Newtonian five-body problem has finitely many spatial central configurations, apart from some explicitly given possible exceptional cases for certain values of the mass parameters.

It seems likely that the spatial five-body central configurations are always finite for positive masses. Our possible exceptional cases often involve some equality between masses, but it is now known that there are finitely many central configurations in the case of five equal masses [10]. Presumably the failure of our methods to resolve our exceptional cases merely reflects the existence of additional syzygies in the leading terms of the generators of our ideal.

8. ACKNOWLEDGMENTS

Some of our computations were run on one of the Sage Foundation's 24-core Sun X4450s, supported by National Science Foundation Grant No. DMS-0821725. We were also both supported by the American Institute of Mathematics. Furthermore, the second author was supported by the German Research Foundation (Deutsche Forschungsgemeinschaft (DFG)) through the Institutional Strategy of the University of Göttingen.

REFERENCES

1. A. Albouy and A. Chenciner, *Le problème des n corps et les distances mutuelles*, *Inv. Math.*, **131** (1998) 151–184.
2. A. Albouy and V. Kaloshin, personal communication.
3. O. Dziobek, *Über einen merkwürdigen Fall des Vielkörperproblems*, *Astron. Nach.* **152** (1900) 33–46.
4. L. Euler, *De motu rectilineo trium corporum se mutuo attrahentium*, *Novi Comm. Acad. Sci. Imp. Petrop.* **11** (1767) 144–151.
5. M. Hampton and R. Moeckel, *Finiteness of relative equilibria of the four-body problem*, *Inventiones Mathematicae* **163** (2006), 289–312.
6. K. Hept and T. Theobald, *Tropical bases by regular projections*, *Proc. Amer. Math. Soc.* **137** (2009), 2233–2241.
7. A. N. Jensen, *Algorithmic Aspects of Gröbner Fans and Tropical Varieties*, Ph.D. Thesis, Department of Mathematical Sciences, University of Aarhus, Denmark (2007).
8. A. N. Jensen, *Gfan, a software system for Gröbner fans*, <http://www.math.tu-berlin.de/~jensen/software/gfan/gfan.html>.
9. J. L. Lagrange, *Essai sur le problème des trois corps*, *Œuvres*, vol. 6 (1772).
10. T.-L. Lee and M. Santoprete, *Central configurations of the five-body problem with equal masses*, *Cel. Mech. and Dynam. Astron.* **104** (2009), 369–381.
11. R. Moeckel, *Generic Finiteness for Dziobek Configurations*, *Trans. Amer. Math. Soc.* **353** (2001) 4673–4686.
12. R. Moeckel, *On central configurations*, *Math. Zeit.* **205** (1990) 499–517.
13. F. R. Moulton, *The Straight Line Solutions of the Problem of n Bodies*, *Ann. of Math.* **12** (1910) 1–17.
14. I. Newton, *Philosophi naturalis principia mathematica*, Royal Society, London (1687).
15. G. Roberts, *A continuum of relative equilibria in the five-body problem*, *Phys. D* **127** (1999) no. 3–4, 141–145.
16. G. Roberts, personal communication.
17. W. A. Stein et al., *Sage: Open Source Mathematical Software (Version 4.3.3)*, The Sage Development Team, 2010, <http://www.sagemath.org>.
18. G.-M. Greuel, G. Pfister, and H. Schönemann: *SINGULAR 3.0 - A Computer Algebra System for Polynomial Computations* in M. Kerber and M. Kohlhase: *Symbolic Computation and Automated Reasoning*, The Calculemus-2000 Symposium. (2001) 227–233.
19. B. Sturmfels, *Gröbner bases and convex polytopes*, volume 8 of *University Lecture Series*, American Mathematical Society, Providence, RI (1996).
20. B. Sturmfels, *Solving systems of polynomial equations*, volume 97 of *CBMS Regional Conference Series in Mathematics*. Published for the Conference Board of the Mathematical Sciences, Washington, DC (2002).
21. C. Witty, *Isolate real roots of real polynomials*, http://www.sagemath.org/doc/reference/sage/rings/polynomial/real_roots.html.

MARSHALL HAMPTON, DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF MINNESOTA, DULUTH, 1117 UNIVERSITY DRIVE, DULUTH, MN, 55812, USA

ANDERS JENSEN, MATHEMATISCHES INSTITUT, GEORG-AUGUST-UNIVERSITÄT
GÖTTINGEN, BUNSENSTR. 3-5, D-37073 GÖTTINGEN GERMANY